


Mice will help reveal the roles of human brown fat

June 2 2017



Adult mice have metabolically active, supraclavicular brown fat depots that share similarities with those in humans. Credit: Miao-Hsueh Chen, Baylor College of Medicine

When it gets cold around you, your body turns up the heat to maintain its normal temperature. The heat is produced by brown adipose tissue, or brown fat, which also plays a role in how the body uses glucose and fat.

However, scientists do not completely understand how brown fat carries out its functions both in health and disease, in part because of the lack of an appropriate animal model. In a paper published in the *Journal of Clinical Investigation Insight*, a team of researchers from several institutions, including Baylor College of Medicine, has filled this gap with the discovery that mice also have brown fat deposits similar to the largest depot found in people. The discovery opens the door to research that might lead to new ways of using brown fat to treat metabolic conditions such as obesity and type 2 diabetes in the future.

"In addition to white adipose tissue, or [white fat](#), people have brown fat, an important contributor to the body's energy balance via the generation of body heat and the participation in metabolic processes," said senior author Dr. Miao-Hsueh Chen, assistant professor of pediatrics and nutrition at Baylor College of Medicine and the USDA/ARS Children's Nutrition Research Center at Baylor and Texas Children's Hospital.

Brown fat contains adipocytes, cells that are rich in small fat-filled droplets and in energy-producing structures called mitochondria. Brown fat adipocytes use fat and glucose as sources of energy. In [mice](#), brown fat activated to produce heat markedly affects the energy balance. For instance, mice housed at temperatures below their normal body temperature (20-22 degrees Celsius) would need to consume 60 percent more food to maintain their normal temperature than mice housed at 30 C. Other experiments have shown that when brown fat is dysfunctional or absent, mice decrease their energy expenditure and become obese.

A mouse model of human brown fat

Studies have indicated that most brown fat in mice is on the back, between the shoulder blades. In people, however, the main depots of brown fat are located above the collar bones and deep in the neck. Scientists think that what they learn by studying mouse brown fat might

not be applicable to people because mouse and human brown fat are at different locations. In the search for a better mouse model, Chen and her colleagues analyzed mouse embryos and found brown fat surrounded by muscles in the neck, including a brown fat depot located above the collar bones, the same location of main human brown fat that had not been described before.

"Further studies showed that adult mice also have brown fat above the collar bones," Chen said. "This is important because studies will be carried out mostly in adults. In addition, mouse brown fat in the [collar bone](#) is morphologically similar to human brown fat in the same location, produces compounds involved in the production of heat and expresses genes similar to those expressed by human brown fat."

Mouse brown fat can change metabolism

In collaboration with Dr. Kristin Stanford, assistant professor of physiology and cell biology at Ohio State University Wexner Medical Center, the research team carried out brown fat transplantation experiments in mice.

"For several years, I've been interested in how to combat obesity and improve metabolic health," Stanford said. "A few years ago, my lab developed a transplantation model looking at the effects of increasing brown fat above the shoulder blade in mice, and we saw a dramatic improvement in metabolic health. When Dr. Chen showed me her data identifying brown fat above the collar bone in mice, I was excited to collaborate and apply our transplantation model."

When the researchers increased the amount of brown fat above the collar bone by transplanting more of it into healthy mice, they saw improvement on the animals' glucose tolerance. This shows that "this brown fat depot, which is remarkably similar to the main brown fat

depot in humans, can be metabolically beneficial," Stanford said. "This study highlights how important this tissue most likely is in humans."

"I am most excited to bring this model to scientists in the field so they can use it to study brown fat," Chen said. "This model is the first step to improve our understanding of the role of human [brown fat](#) in metabolic processes. The [model](#) offers the possibility of carrying out studies that might result in treatments to reverse or prevent diseases such as type 2 diabetes and obesity."

More information: Qianxing Mo et al, Identification and characterization of a supraclavicular brown adipose tissue in mice, *JCI Insight* (2017). [DOI: 10.1172/jci.insight.93166](https://doi.org/10.1172/jci.insight.93166)

Provided by Baylor College of Medicine

Citation: Mice will help reveal the roles of human brown fat (2017, June 2) retrieved 9 April 2024 from <https://medicalxpress.com/news/2017-06-mice-reveal-roles-human-brown.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--