

## Non-shivering thermogenesis can activate metabolic pathways

June 6 2017



(HealthDay)—Non-shivering thermogenesis can transform storage



adipose tissues into metabolically active tissue, according to research published online May 25 in *Obesity Reviews*.

Biff F. Palmer, M.D., from the University of Texas Southwestern Medical Center in Dallas, and Deborah J. Clegg, Ph.D., from Cedars-Sinai Medical Center in Los Angeles, address mechanisms by which nonshivering thermogenesis may offer insight into metabolic pathways that can become targets to facilitate sustainable weight loss.

The researchers note that non-shivering thermogenesis results in activation and expansion of brown and beige adipose tissues; in addition, it activates pathways in skeletal muscle that increase metabolic flux and the activity of muscle fibers via futile calcium cycling across the endoplasmic reticulum. All of these facilitate an increase in metabolism.

"We conclude by emphasizing that non-shivering thermogenesis activates <u>metabolic pathways</u> which could be druggable targets, to facilitate sustainable weight loss," the authors write.

## More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2017 HealthDay. All rights reserved.

Citation: Non-shivering thermogenesis can activate metabolic pathways (2017, June 6) retrieved 3 May 2024 from https://medicalxpress.com/news/2017-06-non-shivering-thermogenesis-metabolic-pathways.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.