

The cost of opioid use during pregnancy

June 14 2017

A new study published today by the scientific journal *Addiction* reveals that the incidence of neonatal abstinence syndrome - often caused by mothers using opioids during pregnancy - is increasing in the United States, and carries an enormous burden in terms of hospital days and costs. The number of US hospital admissions involving neonatal abstinence syndrome increased more than fourfold between the years 2003 and 2012. In 2012, neonatal abstinence syndrome cost nearly \$316 million in the United States.

Neonatal abstinence syndrome (NAS) is a constellation of symptoms that occur in newborn infants exposed to addictive illegal or prescription drugs in utero. Infants affected by NAS typically show a number of neurological symptoms and behaviors (e.g., tremors, seizures) as well as poor feeding and gastrointestinal dysfunction. Standard management of NAS involves the administration of opioids for opioid withdrawal, with additional medications for stubborn cases or instances of multi-drug exposure. This drug administration has been performed traditionally in the hospital setting, consuming valuable and finite hospital resources.

In this study, researchers measured the inflation-adjusted health care provider costs and length of [hospital stay](#) for almost 28,000 infants with NAS, compared with over 3 million infants without NAS. They found that between 2003 and 2012, NAS admissions increased more than fourfold, resulting in a surge in annual costs from \$61 million and 67,869 hospital days in 2003 to nearly \$316 million and 291,168 hospital days in 2012. For an infant affected by NAS, the hospital stay was nearly 3.5 times as long (16.57 hospital days compared with 4.98 days for a non-

NAS patient) and the costs more than three times greater (\$16,893 compared with \$5,610 for a non-affected infant).

In an era of increasingly constrained health-care resources, the rising incidence of NAS has significant implications for policymakers, [hospital](#) administrators and [health care](#) providers nation-wide. Increased public health initiatives are needed to target, educate and provide resources for women of reproductive age to decrease in utero drug exposure. Additionally, given the large variation in screening and treatment of NAS, further study is needed to determine the most effective assessment and management strategies, with a focus on developing therapeutic plans that offer more cost-efficient management of NAS.

More information: Tammy E. Corr et al, The economic burden of neonatal abstinence syndrome in the United States, *Addiction* (2017). [DOI: 10.1111/add.13842](https://doi.org/10.1111/add.13842)

Provided by Society for the Study of Addiction

Citation: The cost of opioid use during pregnancy (2017, June 14) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-06-opioid-pregnancy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--