

Valproic acid concentration drops with ketogenic diet

June 22 2017



(HealthDay)—For patients with epilepsy, anti-epileptic drug serum

concentrations remain stable during the ketogenic diet (KD), apart from those of valproic acid, according to a study published online June 18 in the *Journal of Clinical Pharmacy and Therapeutics*.

G. Heo, Pharm.D., from Yonsei University in Incheon, South Korea, and colleagues examined the potential effects of KD and other dietary therapies in patients with epilepsy. Before and during the treatment, the authors assessed the mean [serum concentrations](#) and mean serum concentrations per weight per daily dose per bioavailability of anti-epileptic drugs. Data were included for 139 patients (81 males; median age, 2.91 years).

The researchers found that the serum concentrations of carbamazepine, lamotrigine, levetiracetam, topiramate, and [valproic acid](#) decreased after the dietary therapies, while there was a slight increase in phenobarbital. The difference was only statistically significant for valproic acid (P

"Most anti-epileptic drug serum concentrations remained stable during the KD and other related dietary therapies except those of valproic acid," the authors write. "Therefore, serum concentrations of valproic acid should be monitored when the KD and other dietary therapies are concomitantly administered."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Valproic acid concentration drops with ketogenic diet (2017, June 22) retrieved 26 April 2024 from <https://medicalxpress.com/news/2017-06-valproic-acid-ketogenic-diet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.