

Adrenal gland activity change under severe stress causing abnormal release of glucocorticoid stress

July 17 2017

Published in the *Proceedings of the National Academy of Sciences* (*PNAS*), the study aimed to find out what the most important factors are to allow fast and flexible responses to stress, and how these factors interact with one another.

Researchers from Bristol's Henry Wellcome Laboratories for Integrative Neuroscience and Endocrinology, in collaboration with colleagues from the University of Exeter's Wellcome Trust Centre for Biomedical Modelling and Analysis, developed a novel mathematical model of the molecular network controlling glucocorticoid synthesis, used the model to predict adrenal responses to <u>stress</u>, and tested these prediction experimentally in vivo in the rat.

Using this interdisciplinary approach, the authors show how the molecular network in the adrenal gland acts dynamically in unstressed physiological conditions, and how the dynamics of these processes change in the presence of <u>inflammatory stimuli</u>, such as infection or during surgery.

Dr Francesca Spiga, one of the lead authors of the study, said: "This is the first study to show just how dynamically complex the adrenal gland <u>response</u> to stress is, and how sensitive is to clinically important perturbations, such as pro-inflammatory cytokines. Our hope is that a better understanding of this system will improve treatment of patients



with inflammatory conditions, such as those undergoing major surgery."

Professor John Terry, from the University of Exeter's Living Systems Institute and one of the senior authors of the study said: "It has long been a mystery whether the <u>adrenal glands</u> secreted glucocorticoids purely under instruction from the brain or whether the gland itself played a role in governing the level of hormones."

"Our latest findings add to a growing body of evidence that stress, and the body's response to stress, is not all in the head, but that the adrenal gland is playing an important role in regulating our <u>stress response</u>."

The findings of this study are important as they show how the dynamics of glucocorticoid production can change under different conditions, which may be responsible for the disrupted rhythms of glucocorticoid hormones associated with disease and ageing.

More information: Francesca Spiga el al., "Dynamic responses of the adrenal steroidogenic regulatory network," *PNAS* (2017). <u>www.pnas.org/cgi/doi/10.1073/pnas.1703779114</u>

Provided by University of Bristol

Citation: Adrenal gland activity change under severe stress causing abnormal release of glucocorticoid stress (2017, July 17) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2017-07-adrenal-gland-severe-stress-abnormal.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.