

Antiplatelet drugs are often inappropriately prescribed in older patients

July 19 2017

A study has found that antiplatelet drugs, such as aspirin, are often inappropriately prescribed in acutely hospitalized older people.

Antiplatelet drugs are typically used to prevent conditions such as heart attacks and strokes, both for individuals at high risk (primary prevention) and for those who have already experienced one of these events (secondary prevention).

In the *British Journal of Clinical Pharmacology* analysis of nearly 2200 patients in Italian and Spanish internal medicine and geriatric clinical wards, 959 (44%) were prescribed an antiplatelet drug. Among patients prescribed for [primary prevention](#), half of them were inappropriately prescribed (52%), being mainly overprescribed (155/209 patients, 74%). On the other hand, a high rate of inappropriate underprescription was also seen in the context of [secondary prevention](#) (222/726 patients, 31%).

More information: *British Journal of Clinical Pharmacology* (2017).
[DOI: 10.1111/bcp.13364](https://doi.org/10.1111/bcp.13364)

Provided by Wiley

Citation: Antiplatelet drugs are often inappropriately prescribed in older patients (2017, July 19)
retrieved 10 April 2024 from

<https://medicalxpress.com/news/2017-07-antiplatelet-drugs-inappropriately-older-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.