

Babies with hearing loss need early intervention, but only half get it

July 13 2017, by Lisa Marshall



An infant has her hearing screened. Credit: Christine Yoshinaga-Itano

Children with hearing loss who are diagnosed by 3 months of age and receive interventions by 6 months develop a far greater vocabulary than those whose diagnosis and treatment come later, according to a CU Boulder study published this week in the journal *Pediatrics*. Yet 17 years after early detection guidelines were established, nearly half of babies with hearing loss aren't meeting them, the study found.

"We still have some work to do," said lead author Christine Yoshinaga-

Itano, an audiologist and research professor in the Institute of Cognitive Science. "Because the brain is so pliable in those early [months](#), the sooner we can get them diagnosed and get them access to language, the more likely they will be able to develop on track with their peers."

Funded by the Centers for Disease Control, the paper is the first multistate study to assess the impact of Early Hearing Detection Intervention (EHDI) 1-3-6 guidelines. The Joint Committee on Infant Hearing, founded by late CU Professor Marion Downs, established the guidelines in 2000. They recommend that all newborns be screened for hearing loss within one month, those who test positive be evaluated by a specialist within three months, and those diagnosed with hearing loss begin interventions (such as family support services, listening and spoken language intervention, sign language instruction, cochlear implants or hearing aids) within six months.

Today, 96 percent of U.S. infants are screened by one month. But many with hearing loss do not meet steps two and three, often due to family difficulties getting transportation, appointments, time off, or funding for [hearing aids](#), which are not always covered by insurance. Navigating the complex U.S. healthcare system can also be challenging, notes Yoshinaga-Itano.

The study looked at 448 [children](#) with hearing loss in both ears, ages 8 months to 39 months, across 12 states. Only 58 percent had met the EHDI 1-3-6 guidelines.

Researchers also measured the number of words the children used either in spoken or sign language at different ages and compared it to hearing peers. (On average, hearing children have about 600 words in their expressive vocabulary by 30 months.)

A Vocabulary Quotient (VQ) score on par with a hearing peer would be

100. Children with hearing loss who had met the guidelines and had no additional disabilities scored 82 on average, the low end of normal range. Those who had not met the guidelines scored below 70, ranking in the lowest 10th percentile.

The [study](#) was a national follow-up to Yoshinaga-Itano's landmark 1998 study in *Pediatrics*, which looked only at children with hearing loss in Colorado, a national leader in early intervention. It similarly found that while those who were diagnosed and received interventions by six months developed language on par with their peers, those diagnosed later had poorer outcomes.

"We showed that failure to diagnose [hearing loss](#) early can create an environmentally induced and preventable secondary disability, making children function much like children with cognitive delay," she said. That study helped kick-start national universal newborn hearing screening efforts.

The new study also found that children who have more hearing at birth, parents with a better education, or parents who are deaf (and already familiar with sign language or other means of communicating) also have better vocabulary scores.

But those variables are harder to change, Yoshinaga-Itano says.

"We can't change how much hearing a child has at birth or the educational background of a parent, but we can develop better systems," she says. "Policymakers need to do whatever they can to make transitions from one step to another as seamless as possible so parents can meet the 1,3,6. And parents should know that there is an urgency to assuring that children who are deaf or hard of [hearing](#) have access to [language](#) as quickly as possible."

More information: Christine Yoshinaga-Itano et al, Early Hearing Detection and Vocabulary of Children With Hearing Loss, *Pediatrics* (2017). [DOI: 10.1542/peds.2016-2964](https://doi.org/10.1542/peds.2016-2964)

Provided by University of Colorado at Boulder

Citation: Babies with hearing loss need early intervention, but only half get it (2017, July 13)
retrieved 28 April 2024 from
<https://medicalxpress.com/news/2017-07-babies-loss-early-intervention.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--