


Co-infection with two common gut pathogens worsens malnutrition in mice

July 27 2017



Histopathology of duodenum in infected mice fed PD diet (20x, H&E, arrows designate trophozoites). Credit: Bartelt LA, *et al.* (2017)

Two gut pathogens commonly found in malnourished children combine to worsen malnutrition and impair growth in laboratory mice, according to new research published in *PLOS Pathogens*.

Malnourished children often face infection with pathogenic microbes that colonize the intestines. These infections disrupt healthy gut microbial communities and harm metabolism and immune system function, worsening [malnutrition](#) and impairing children's growth and development. However, the precise effects of co-infection with multiple [pathogens](#) in [malnourished children](#) are poorly understood.

To gain new insights, Luther Bartelt of the University of North Carolina at Chapel Hill and colleagues developed a new laboratory mouse model of co-infection during malnutrition. They fed weaned mice a protein-deficient diet and infected them with *Giardia lamblia* and, two weeks later, *enteroaggregative Escherichia coli* (EAEC)—two of the pathogens most commonly found in malnourished children.

The researchers then used an array of analytical tools—including stool and urine analysis, flow cytometry, light microscopy, [nuclear magnetic resonance](#) spectroscopy, and 16S rRNA analysis—to observe the wide-ranging effects of this sequential co-infection.

They found that *Giardia* and EAEC combined to increase weight loss in the young mice. This appeared to be a result of both impaired metabolism and worsened [immune system function](#) in the mucosal lining of the mice's intestines. Co-infection also amplified protein breakdown by gut microbes and simultaneously interfered with the ability for the

mouse metabolism to adapt to protein deficiency.

These findings and future studies in similar co-infection mouse models could help reveal new insights that cannot be gleaned from single-pathogen studies alone. They could also help inform ongoing long-term studies of malnutrition in children and, ultimately, the development of new treatments or better combinations of existing treatments to restore healthy gut conditions in malnourished children.

"We observed unique inflammatory and metabonomic consequences of *Giardia* and EAEC infections, not only separately, but also during co-infection," the authors explain. "These findings inform our understanding of similar perturbations seen in malnourished [children](#), and change how we think about mechanisms driving multi-enteropathogen-associated enteropathy."

More information: Bartelt LA, Bolick DT, Mayneris-Perxachs J, Kolling GL, Medlock GL, Zaenker EI, et al. (2017) Cross-modulation of pathogen-specific pathways enhances malnutrition during enteric co-infection with *Giardia lamblia* and enteroaggregative *Escherichia coli*. *PLoS Pathog* 13(7): e1006471. doi.org/10.1371/journal.ppat.1006471

Provided by Public Library of Science

Citation: Co-infection with two common gut pathogens worsens malnutrition in mice (2017, July 27) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-07-co-infection-common-gut-pathogens-worsens.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
