

## **Common strength 'genes' identified for first time**

July 12 2017



Skeletal muscle tissue. Credit: University of Michigan Medical School

Common genetic factors that influence muscle strength in humans have been identified for the first time in a study led by researchers from the University of Cambridge and published today in *Nature Communications*.

The researchers used data on hand grip <u>strength</u> from more than 140,000 participants in the UK Biobank study, combined with 50,000 additional individuals from the UK, Netherlands, Denmark and Australia, to identify sixteen common genetic variants that are associated with <u>muscle</u> strength.



Dan Wright, joint first author on this paper and a PhD student at the Medical Research Council Epidemiology (MRC) Unit at the University of Cambridge, said: "The very large number of individuals participating in UK Biobank provides a powerful resource for identifying genes involved in complex traits such as muscle strength, and helps us understand their underlying biology and its relevance to health."

Many of these variants were located within or near to genes known to play a role in biological processes highly relevant to muscle function, including the structure and function of <u>muscle fibres</u>, and the communication of the nervous system with muscle cells.

Mutations in some of the genes highlighted are also known to be associated with severe monogenic syndromes - conditions caused by a single genetic mutations - characterised by compromised <u>muscle</u> <u>function</u>. This demonstrates that <u>genetic variation</u> in genes which cause serious muscular conditions may also influence differences in strength in the general population.

Dr Robert Scott, who co-led the study with colleagues from the MRC Epidemiology Unit, said: "While we have long suspected a role for genetics in the variation in muscle strength, these findings give the first insights into some of the specific genetic variants that underpin variation in strength.

"These could be important steps towards identifying new treatments to prevent or treat muscle weakness."

Hand grip strength has been reported to be associated with many health outcomes, including risk of mortality, cardiovascular disease, and fracture - although it has been unclear whether variation in strength actually causes these outcomes, or simply reflects underlying disease processes.



Using the sixteen genetic variants identified for strength, the researchers were able to investigate the hypothesised causal link between strength and these adverse health outcomes. Their study found no evidence that lower strength causally increases risk of death or cardiovascular disease, but they did find evidence that higher muscular strength reduces risk of fracture, supporting the use of strength training interventions as a strategy to reduce risk of fractures.

Professor Nick Wareham, director of the MRC Epidemiology Unit and a senior author of the study, noted: "This work highlights the importance of <u>muscle strength</u> in the prevention of fractures and the complications which can often follow a fall."

**More information:** Willems, SM et al. Large-scale GWAS identifies multiple loci for hand grip strength providing biological insights into muscular fitness. *Nature Communications*; 12 July 2017; DOI: 10.1038/ncomms16015

## Provided by University of Cambridge

Citation: Common strength 'genes' identified for first time (2017, July 12) retrieved 23 May 2024 from <u>https://medicalxpress.com/news/2017-07-common-strength-genes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.