

Sleep disorders may increase cognitive problems particularly in those at risk for Alzheimer's

July 21 2017



Sleep disorders may affect attention, memory in patients at risk for Alzheimer's disease. Credit: ATS

People who carry a genetic susceptibility to Alzheimer's disease appear to be at greater risk of diminished cognition from sleep-disordered breathing than those without the susceptibility, according to new research published online, ahead of print in the *Annals of the American Thoracic Society*.

In "Greater Cognitive Deficits with Sleep-Disordered Breathing among Individuals with Genetic Susceptibility to Alzheimer's Disease: The Multi-Ethnic Study of Atherosclerosis," researchers report that study participants carrying the apolipoprotein ϵ -4 (APOE- ϵ 4) allele showed greater cognitive deficits with the various indices of sleep-disordered breathing compared to those without the allele.

APOE is a major cholesterol carrier that supports injury repair in the brain. Other studies have shown that those carrying the alternate form of the gene, ϵ 4 allele, are at increased risk of Alzheimer's disease. Estimates are that 20 percent of the population carries the ϵ 4 allele.

"Previous studies have shown inconsistent findings between sleep-disordered breathing and cognition, which may be due to the different tests used," said lead study author Dayna A. Johnson, PhD, MPH, MS, MSW, instructor of medicine at Brigham and Women's Hospital and Harvard Medical School.

Dr. Johnson and colleagues investigated the association in a diverse sample using several indicators of sleep-disordered breathing and cognition. They also evaluated whether the presence of the APOE- ϵ 4 allele, which is known to increase risk of Alzheimer's disease, influenced the link between sleep-disordered breathing and cognition.

The authors analyzed data from 1,752 participants (average age 68) in the Multi-Ethnic Study of Atherosclerosis (MESA) who underwent an in-home polysomnography (sleep) study, completed standardized sleep

questions, and a battery of tests to measure their cognition. The authors defined sleep-disordered breathing as an apnea-hypopnea index (AHI), which measures the number of stopped or shallow breaths per hour, as $AHI > 15$, and sleep apnea syndrome as $AHI > 5$ (below 5 is normal) plus self-reported sleepiness (based on a standardized scale).

The study found:

- Increased overnight hypoxemia (oxygen saturation below 90 percent) or increased [daytime sleepiness](#) was associated with poorer attention and memory.
- More daytime sleepiness was also associated with slower cognitive processing speed.
- Sleep apnea syndrome was associated with poorer attention and processing speed.
- These associations were strongest in APOE-ε4 carriers.

The researchers adjusted for race, age, body mass index, education level, smoking status, hypertension, diabetes, benzodiazepine use, and depressive symptoms.

Dr. Johnson said that, overall, the effects of the various sleep factors they measured on cognition were small, but in the range previously reported for several other lifestyle and health risk factors for dementia. Screening and treating sleep-disordered breathing, she added, may help reduce a person's risk of dementia, especially if that individual carries APOE-ε4.

"Our study provides further evidence that sleep-disordered breathing negatively affects attention, processing speed and memory, which are robust predictors of cognitive decline," said senior study author Susan Redline, MD, MPH, Peter C. Farrell Professor of Sleep Medicine, Harvard Medical School.

"Given the lack of effective treatment for Alzheimer's disease, our results support the potential for sleep-disordered [breathing](#) screening and treatment as part of a strategy to reduce dementia risk."

More information: "Genetic susceptibility to #Alzheimer's increases chance that poor #sleep will impair attention, memory and cognitive processing," *Annals of the American Thoracic Society*, 2017.

Provided by American Thoracic Society

Citation: Sleep disorders may increase cognitive problems particularly in those at risk for Alzheimer's (2017, July 21) retrieved 11 July 2024 from <https://medicalxpress.com/news/2017-07-disorders-cognitive-problems-alzheimer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--