

# EU report: More evidence on link between antibiotic use and antibiotic resistance

July 27 2017

---



Staphylococcus aureus - Antibiotics Test plate. Credit: CDC

The European Food Safety Authority, the European Medicines Agency and the European Centre for Disease Prevention and Control are

concerned about the impact of use of antibiotics on the increase in antibiotic-resistant bacteria. The [report](#) presents new data on antibiotic consumption and antibiotic resistance and reflects improved surveillance across Europe.

Vytė Andriukaitis, European Commissioner for Health and Food Safety says:

"To contain [antibiotic resistance](#) we need to fight on three fronts at the same time: human, animal and the environment. This is exactly what we are trying to achieve in the EU and globally with our recently launched EU Action Plan on [antimicrobial resistance](#). This new report confirms the link between antibiotic consumption and antibiotic resistance in both humans and food-producing animals."

## A diverse picture

Experts highlight that there are still important differences across the EU in the use of antibiotics in animals and humans. Cutting their unnecessary use will have an impact on the occurrence of resistance.

Overall antibiotic use is higher in food-producing animals than in humans, but the situation varies across countries and classes of antibiotics. In particular, a class of antibiotics called polymyxins (which includes colistin) is used widely in the veterinary sector. It is also increasingly used in hospitals to treat multidrug-resistant infections. Other antibiotics are more often used in humans than in animals. These include third- and fourth-generation cephalosporins and quinolones, antibiotics that are also considered critically important for [human](#) health.

## Links between antibiotic use and resistance

Experts note that resistance to quinolones, used to treat salmonellosis and campylobacteriosis in humans, is associated with use of these antibiotics in animals. The use of third- and fourth-generation cephalosporins for the treatment of infections caused by *E. coli* and other bacteria in humans is associated with resistance to these antibiotics in *E. coli* found in humans.

## Cooperation spurs better reporting and analysis

The report is the result of close cooperation between the three EU agencies, each drawing on their specific expertise and data from monitoring of antibiotic resistance and antibiotic consumption in [animals](#) and humans.

The conclusions are in line with those of the [first report published by the three agencies in 2015](#) . However, the availability of better quality data allowed a more sophisticated analysis.

Experts of the three agencies recommend further research to understand better how use of [antibiotics](#) and resistance affect one another.

Provided by European Centre for Disease Prevention and Control (ECDC)

Citation: EU report: More evidence on link between antibiotic use and antibiotic resistance (2017, July 27) retrieved 10 May 2024 from <https://medicalxpress.com/news/2017-07-eu-evidence-link-antibiotic-resistance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---