

False-positive mammogram results linked to spike in anxiety prescriptions

July 6 2017



Credit: Getty Images

Women who experience a false-positive mammogram result are more likely to begin medication for anxiety or depression than women who received an immediate negative result, according to a study led by Penn

State researcher Joel Segel. The finding highlights the importance of swift and accurate follow-up testing to rule out a breast cancer diagnosis.

The study found that patients who receive a false-positive mammogram result are also prescribed anxiety or depression [medication](#) at a rate 10 to 20 percent higher than patients who receive an immediate negative result. These prescriptions are new and not continuations of previously prescribed medicines.

A false-positive result is one where a suspicious finding on the [screening](#) mammogram leads to additional testing that does not end up leading to a breast cancer diagnosis.

Additionally, within that group of patients who required more than one test to resolve the false-positive there was a 20 to 30 percent increase in those beginning to take anxiety or depression medications. The increase was particularly noticeable among [women](#) with commercial insurance who required multiple tests to rule out a breast cancer diagnosis.

"The results suggest that efforts to quickly resolve initially positive findings including same-day follow-up tests may help reduce anxiety and even prevent initiation of anxiety or depression medication," said Segel, assistant professor of health policy and administration at Penn State.

This study demonstrates that some women who experience a false-positive mammogram may need additional follow-up care to effectively handle the increased anxiety that may accompany the experience, Segel said.

More importantly, from a practitioner standpoint, the study identifies sub-populations who may be most at risk of increased anxiety following a false-positive mammogram, Segel said. Specifically, women whose false-positive result requires more than one follow-up test to resolve,

women with commercial insurance who undergo a biopsy, women who wait longer than one week to receive a negative result, and women who are under age 50 may all be at higher risk of experiencing clinically significant anxiety or depression.

"Regular breast cancer screening is critical to early detection," Segel said. "Patients should continue to work with their providers to ensure they are receiving guideline-appropriate screening and should follow up with their providers if they experience either anxiety or depression following screening or any type of care."

Researchers studied commercial- and Medicaid-claims databases to identify women ages 40 to 64 who underwent screening mammography with no prior claims for [anxiety](#) or [depression](#) medications.

The findings recently appeared in *Medical Care*.

More information: Joel E. Segel et al. The Effect of False-positive Mammograms on Antidepressant and Anxiolytic Initiation, *Medical Care* (2017). [DOI: 10.1097/MLR.0000000000000760](https://doi.org/10.1097/MLR.0000000000000760)

Provided by Pennsylvania State University

Citation: False-positive mammogram results linked to spike in anxiety prescriptions (2017, July 6) retrieved 24 April 2024 from <https://medicalxpress.com/news/2017-07-false-positive-mammogram-results-linked-spike.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|