


Girl's HIV infection seems under control without AIDS drugs

July 24 2017, by Marilyn Marchione



This undated photo provided by the Centers for Disease Control and Prevention shows a scanning electron micrograph of multiple round bumps of the HIV-1 virus on a cell surface. In a report released on Monday, July 24, 2017, researchers said a South African girl born with the AIDS virus has kept her infection suppressed for 8 1/2 years after stopping anti-HIV medicines—more evidence that early treatment can occasionally cause a long remission that, if it lasts, would be a form of cure. (Cynthia Goldsmith/Centers for Disease Control and Prevention via AP)

A South African girl born with the AIDS virus has kept her infection suppressed for more than eight years after stopping anti-HIV medicines—more evidence that early treatment can occasionally cause a long remission that, if it lasts, would be a form of cure.

Her case was revealed Monday at an AIDS conference in Paris, where researchers also gave encouraging results from tests of shots every month or two instead of daily pills to treat HIV.

"That's very promising" to help people stay on [treatment](#), the U.S.'s top AIDS scientist, Dr. Anthony Fauci, said of the prospects for long-acting drugs.

Current treatments keep HIV under control but must be taken lifelong. Only one person is thought to be cured—the so-called Berlin patient, a man who had a bone marrow transplant in 2007 from a donor with [natural resistance](#) to HIV.

But transplants are risky and impractical to try to cure the millions already infected. So some researchers have been aiming for the next best thing—long-term remission, when the immune system can control HIV without drugs even if signs of the virus remain.

Aggressive treatment soon after infection might enable that in some cases, and the South African girl is the third child who achieved a long remission after that approach.

She was in a study sponsored by the agency Fauci heads, the National Institute of Allergy and Infectious Diseases, that previously found that early versus delayed treatment helped babies survive.

The girl, who researchers did not identify, started on HIV drugs when she was 2 months old and stopped 40 weeks later. Tests when she was 9

1/2 years old found signs of virus in a small number of immune system cells, but none capable of reproducing. The girl does not have a gene mutation that gives natural resistance to HIV infection, Fauci said, so her remission seems likely due to the early treatment.

The previous cases:

—A French teen who was born with HIV and is now around 20 has had her infection under control despite no HIV medicines since she was roughly 6 years old.

—A Mississippi baby born with HIV in 2010 suppressed her infection for 27 months after stopping treatment before it reappeared in her blood. She was able to get the virus under control again after treatment resumed.

At least a dozen adults also have had remissions lasting for years after stopping HIV medicines.

A study underway now is testing whether treating HIV-infected newborns within two days of birth can control the virus later after treatment stops. It started in 2014 in South America, Haiti, Africa and the United States, and some of the earliest participants might be able to try stopping treatment later this year.

Treatment might get easier if two large studies underway now confirm results reported Monday from a study testing a long-acting combo of two HIV drugs—Janssen's rilpivirine and ViiV Healthcare's cabotegravir.

Cabotegravir is experimental; rilpivirine is sold now as Edurant and used in combination with other drugs for treating certain types of HIV patients.

After initial treatment to get their virus under control, about 300 study participants were given either daily combination therapy pills or a shot every four or eight weeks of the long-acting drug duo to maintain control.

After nearly two years, 94 percent on eight-week shots, 87 percent on four-week shots and 84 percent on daily pills had their infections suppressed, with similar rates of side effects.

"The results were good regardless of whether people came monthly or every two months for their treatment. This has important policy implications," said Dr. Linda-Gail Bekker, deputy director of the Desmond Tutu HIV Centre at the University of Cape Town in South Africa, and a co-leader of the conference.

The study was sponsored by the drugmakers. Results were published in the British medical journal *Lancet*.

Two large studies aimed at winning approval to sell the treatment are testing the monthly shots. Janssen said in a statement that good results from eight-week shots warrant reconsidering the longer approach.

If it works, "this will have a huge impact on how we manage that very important group of people who are not able to access and take drugs on a day-to-day basis," such as those with mental health or [drug](#) abuse problems, said Dr. Steven Deeks, an AIDS specialist at the University of California, San Francisco.

More information: Reference: A Violari et al. Viral and host characteristics of a child with perinatal HIV-1 following a prolonged period after ART cessation in the CHER trial. 9th IAS Conference on HIV Science, Paris (2017).

© 2017 The Associated Press. All rights reserved.

Citation: Girl's HIV infection seems under control without AIDS drugs (2017, July 24) retrieved 25 April 2024 from <https://medicalxpress.com/news/2017-07-girl-hiv-infection-aids-drugs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.