

# The hidden chemicals in hookah tobacco smoke

July 18 2017, by Katie White

---



Smoking hookah has gained popularity in recent years. Credit: San Diego State University

Thanks to its sweet smell and taste, hookah has gained popularity in recent years, especially among teens and young adults. Hookah users

inhale smoke, which is generated by heating hookah tobacco that is fermented with molasses and fruits and combined with burning charcoal. The smoke passes through a partially filled water jar which creates a bubbly sound, making smoking hookah appear more innocent than cigarettes. But first impressions can be deceiving.

A new study out of San Diego State University's Center for Behavioral Epidemiology and Community Health shows that burning [hookah](#) tobacco creates a dangerous byproduct: the chemical compound acrolein. Lead author Nada Kassem, a research scientist in SDSU's Graduate School of Public Health, and her collaborators found that the use of tobacco in a hookah exposes smokers and non-smokers to potentially dangerous levels of the respiratory irritant that is believed to cause cancer and heart disease.

The study was funded by the American Cancer Society and was released in the journal *Nicotine and Tobacco Research*.

In this study, Kassem and her colleagues analyzed levels of 3-HPMA, a metabolite of acrolein, in the urine of 105 hookah smokers, and 103 non-smokers exposed to secondhand smoke at indoor hookah events where hookah tobacco was smoked exclusively. Participants provided urine samples the morning of and the morning after attending the event.

"We found that after spending an average of three hours at a hookah party, the levels of 3-HPMA in urine increased by an average of nearly 1.5 times in both those [smoking](#) hookah and those simply attending the event without smoking hookah," said Kassem. "The acrolein levels were even higher in people who smoked hookah on a daily basis. They had acrolein levels four times higher than those found in a representative sample of non-tobacco users in the United States."

In a previous study, Kassem found that children five years or younger

who live in homes of hookah smokers were exposed to acrolein. For example, children of exclusive daily hookah smokers had 1.9 times higher levels of urinary 3-HPMA than their counterparts who live in non-smokers' homes.

Kassem has been investigating the dangers of hookah smoke for several years and is part of a group of SDSU researchers who investigate the harmful effects of second- and thirdhand smoke. She previously published a study about benzene levels in those exposed to hookah.

Kassem has been awarded a \$1.4 million grant from the National Institutes of Health to further research the dangers of hookah smoke and is currently in the process of collecting more data.

She says all of her current and previous research points to the one conclusion: Smoking hookah is not a safe alternative to smoking other forms of tobacco, and she suggests stricter regulations for hookah bars and events.

"Our results support regulating hookah tobacco products, including reducing additives that produce acrolein, including sugars that sweeten the smell and taste of the tobacco, and humectants, which are used to maintain the moisture of the tobacco," said Kassem. "We also recommend posting health warning signs for indoor smoking in hookah lounges, and encouraging bans of indoor hookah [tobacco](#) smoking in private homes."

**More information:** Nada O.F. Kassem et al. Acrolein exposure in hookah smokers and non-smokers exposed to hookah tobacco Secondhand Smoke: Implications for regulating hookah tobacco products, *Nicotine & Tobacco Research* (2017). [DOI: 10.1093/ntr/ntx133](https://doi.org/10.1093/ntr/ntx133)

Provided by San Diego State University

Citation: The hidden chemicals in hookah tobacco smoke (2017, July 18) retrieved 18 April 2024 from <https://medicalxpress.com/news/2017-07-hidden-chemicals-hookah-tobacco.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.