


House dust spurs growth of fat cells in lab tests

July 12 2017



Credit: CC0 Public Domain

Poor diet and a lack of physical activity are major contributors to the

world's obesity epidemic, but researchers have also identified common environmental pollutants that could play a role. Now one team reports in ACS' journal *Environmental Science & Technology* that small amounts of house dust containing many of these compounds can spur fat cells to accumulate more triglycerides, or fat, in a lab dish.

Endocrine-disrupting chemicals, or EDCs, are synthetic or naturally occurring compounds that can interfere with or mimic the body's hormones. EDCs, such as flame retardants, phthalates and bisphenol-A, are known for their potential effects on reproductive, neurological and immune functions. But animal studies also suggest that early life exposure to some EDCs can cause weight gain later in life, and, as a result, have been called "obesogens." Some manufacturers have reduced the use of EDCs in products, but many are still ubiquitous in consumer goods. And they wind up in indoor dust that can be inhaled, ingested, or absorbed through the skin. The U.S. Environmental Protection Agency estimates that children consume 50 milligrams of house dust each day. Concerned about the potential effects EDCs in dust might have on children's health, Heather Stapleton and colleagues wanted to see if the compounds in house dust might have an effect on [fat cells](#).

The researchers collected samples of indoor dust from 11 homes in North Carolina and tested extracts from the samples in a mouse pre-adipocyte cell model, 3T3-L1 cells, often used to test compounds for potential effects on the accumulation of triglycerides, a type of fat. Extracts from seven of the 11 [dust samples](#) triggered the pre-adipocytes to develop into mature fat cells and accumulate triglycerides. Extracts from nine samples spurred the cells to divide, creating a larger pool of precursor fat cells. Only one dust sample had no effect.

Additionally, among the 44 individual common house dust contaminants tested in this model, pyraclostrobin (a pesticide), the flame-retardant TBPD, and DBP, a commonly used plasticizer, had the strongest fat-

producing effects. This suggests that the mixture of these chemicals in [house](#) dust is promoting the accumulation of triglycerides and fat [cells](#), the researchers say. Amounts of dust as low as 3 micrograms—well below the mass of [dust](#) that children are exposed to daily—caused measurable effects. Thus, the researchers also suggest that [house dust](#) is a likely exposure source of chemicals that may be able to disrupt metabolic health, particularly in children.

More information: "Characterization of Adipogenic Activity of House Dust Extracts and Semi-Volatile Indoor Contaminants in 3T3-L1 Cells" *Environmental Science & Technology* (2017).

pubs.acs.org/doi/abs/10.1021/acs.est.7b01788

Provided by American Chemical Society

Citation: House dust spurs growth of fat cells in lab tests (2017, July 12) retrieved 25 April 2024 from <https://medicalxpress.com/news/2017-07-house-spurs-growth-fat-cells.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--