

# Brief interactions spur lasting waves of gene activity in the brain

July 17 2017

---



A new study measured robust changes in brain gene expression for up to two hours after a three-spined stickleback fish encountered another fish. Credit: L. Brian Stauffer

A five-minute encounter with an outsider spurs a cascade of changes in

gene activity in the brain that can last for hours, researchers report in a study of stickleback fish.

The research, described in the journal *PLOS Genetics*, is one of three recent studies - the others conducted in honey bees and mice - to see waves of changes in [gene expression](#) in the brain 30 minutes to two hours after contact with an intruder.


"We are discovering that social interactions are extremely potent; they provoke big changes in gene [expression](#) in the brain," said University of Illinois animal biology professor Alison Bell, who studies behavior in three-spined [stickleback fish](#). "These very subtle social interactions are getting under the skin and becoming embedded in the brain. Studies like ours are beginning to show how that actually works."

The observed changes included modifications to chromatin accessibility - a selective unpacking of the DNA that allows some genes, but not others, to be transcribed and translated into proteins.

"We used to think that DNA was so tightly packaged that it took a long time for that packaging to become unwrapped," Bell said. "But our study shows that the unpackaging can happen fast - within minutes to hours - and we show that something really subtle, like a social interaction, can provoke dramatic changes in chromatin accessibility."

Bell and her colleagues tracked changes in gene expression 30, 60 and 120 minutes after an encounter. They focused on two parts of the [brain](#) - the telencephalon, which is important to learning and memory; and the diencephalon, which integrates social information and hormonal influences.

"We compared experimental fish - those that had the encounter - with control fish, who remained alone in their tanks," she said.



The study measured changes in the expression of hundreds of genes in two brain regions in the fish: the telencephalon, which is important to learning and memory; and the diencephalon, which integrates social information and hormonal influences. Credit: Graphic by Julie McMahon

Hundreds of genes were expressed differently in the experimental and control fish. They also varied in the expression of numerous transcription factors, which regulate the activity of other genes.

Graduate Student [Syed Abbas Bukhari](#), who is first author on the paper, led an analysis of the functions of groups of genes whose expression patterns paralleled one another at 30, 60 and 120 minutes.

"This revealed that the genes that change in tandem over time have similar functions," Bell said. Such functional groups include genes that contribute to immunity, hormone activity, metabolism or homeostasis.

In the experimental fish, hormone gene expression was highest at 30 minutes, metabolism genes peaked at 60 minutes and [genes](#) linked to immune function and homeostasis were upregulated the most two hours after the encounter.

"We're seeing changes in gene expression even two hours after the fish interacts with an intruder. Clearly, those changes are not about the animal's initial behavior," Bell said. "We haven't studied learning and memory, but we think that these chromatin modifications and changes in gene expression are all about setting up their brains to respond to threats in the future."

Previous studies suggest that the [fish](#) are learning the limits of their territory, Bell said. "Their territory basically ends where they successfully chased away an intruder," she said.

"Our findings made me feel very humble; I really did not appreciate how much is going on in these guys' brains," Bell said. "If you watch sticklebacks in natural populations, they're constantly fighting with their neighbors. Our study just looked at what happened after a five-minute interaction, once. They're doing this all day long, over and over again."

**More information:** Syed Abbas Bukhari et al, Temporal dynamics of neurogenomic plasticity in response to social interactions in male threespined sticklebacks, *PLOS Genetics* (2017). [DOI: 10.1371/journal.pgen.1006840](https://doi.org/10.1371/journal.pgen.1006840)

Provided by University of Illinois at Urbana-Champaign

Citation: Brief interactions spur lasting waves of gene activity in the brain (2017, July 17)  
retrieved 4 May 2024 from

<https://medicalxpress.com/news/2017-07-interactions-spur-gene-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--