

MRSA emerged years before methicillin was even discovered

July 19 2017



A colorized scanning electron micrograph of MRSA. Credit: National Institute of Allergy and Infectious Diseases

Methicillin resistant *Staphylococcus aureus* (MRSA) emerged long before the introduction of the antibiotic methicillin into clinical practice, according to a study published in the open access journal *Genome Biology*. It was the widespread use of earlier antibiotics such as penicillin rather than of methicillin itself which caused MRSA to emerge,

researchers at the University of St Andrews, and the Wellcome Trust Sanger Institute, UK suggest.

The researchers found that *S. aureus* acquired the gene that confers methicillin [resistance](#) - *mecA* - as early as the mid-1940s - fourteen years before the first use of methicillin.

Professor Matthew Holden, molecular microbiologist at the University of St Andrews, the corresponding author said: "Our study provides important lessons for future efforts to combat [antibiotic resistance](#). It shows that new drugs which are introduced to circumvent known [resistance mechanisms](#), as methicillin was in 1959, can be rendered ineffective by unrecognized, pre-existing adaptations in the bacterial population. These adaptations happen because - in response to exposure to earlier [antibiotics](#) - resistant bacterial strains are selected instead of non-resistant ones as bacteria evolve."

The *mecA* gene confers resistance by producing a protein called PBP2a, which decreases the binding efficiency of antibiotics used against *S. aureus* to the bacterial cell wall. The introduction of penicillin in the 1940s led to the selection of *S. aureus* strains that carried the methicillin resistance gene.

Dr. Catriona Harkins, clinical lecturer in dermatology at the University of Dundee, the first author of the study said: "Within a year of methicillin being first introduced to circumvent penicillin resistance, strains of *S. aureus* were found that were already resistant to methicillin. In the years that followed resistance spread rapidly in and outside of the UK. Five decades on from the appearance of the first MRSA, multiple MRSA lineages have emerged which have acquired different variants of the resistance gene."

To uncover the origins of the very first MRSA and to trace its

evolutionary history, the researchers sequenced the genomes of a unique collection of 209 historic *S. aureus* isolates. The oldest of these isolates were identified over 50 years ago by the *S. aureus* reference laboratory of Public Health England and have been stored ever since in their original freeze-dried state. The researchers also found [genes](#) in these isolates that confer resistance to numerous other antibiotics, as well as genes associated with decreased susceptibility to disinfectants.

Professor Holden said: "*S. aureus* has proven to be particularly adept at developing resistance in the face of new antibiotic challenges, rendering many antibiotics ineffective. This remains one of the many challenges in tackling the growing problem of antimicrobial resistance. In order to ensure that future antibiotics retain their effectiveness for as long as possible, it is essential that effective surveillance mechanisms are combined with the use of genome sequencing to scan for the emergence and spread of resistance."

More information: Catriona P. Harkins et al, Methicillin-resistant *Staphylococcus aureus* emerged long before the introduction of methicillin into clinical practice, *Genome Biology* (2017). [DOI: 10.1186/s13059-017-1252-9](#)

Provided by BioMed Central

Citation: MRSA emerged years before methicillin was even discovered (2017, July 19) retrieved 20 March 2024 from <https://medicalxpress.com/news/2017-07-mrsa-emerged-years-methicillin.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--