

# Significantly higher serum melatonin in human myopes

July 28 2017

---



(HealthDay)—Human myopes exhibit significantly higher serum

melatonin (Mel) concentration than non-myopes, according to a study published online July 18 in *Ophthalmic & Physiological Optics*.

Stephanie Kearney, from the University of Ulster in Coleraine, U.K., and colleagues examined the correlation between myopia and [serum](#) concentrations of dopamine (DA) and Mel in 54 participants (aged  $19.1 \pm 0.81$  years); nine participants were lost to follow-up. Solid [phase](#) extraction-liquid chromatography-tandem mass spectrometry was used to measure morning [serum concentrations](#) of DA and Mel in September/October 2014 (phase 1) and March/April 2016 (phase 2). The authors also recorded axial length (AL), corneal radii (CR), and spherical equivalent refraction (SER).

The researchers found that at phases 1 and 2, myopes exhibited significantly higher Mel concentrations than non-myopes (median differences, 10 and 7.3  $\text{pg mL}^{-1}$ , respectively; both  $P < 0.006$ ). There was a positive association for Mel concentrations with more negative SER (all  $r \geq -0.53$ ; all  $P$

"This study reports for the first time in humans that myopes exhibit higher serum Mel concentrations than non-myopes," the authors write. "This may indicate a role for light exposure and circadian rhythm in the human myopic growth mechanism."

**More information:** [Abstract](#)  
[Full Text](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Significantly higher serum melatonin in human myopes (2017, July 28) retrieved 25 April 2024 from <https://medicalxpress.com/news/2017-07-significantly-higher-serum-melatonin-human.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.