

Thyroid cancer tied to regular thyroxine use in hypothyroidism

July 31 2017



(HealthDay)—Patients with primary hypothyroidism who are regular

thyroxine users have increased risk of thyroid cancer, according to a study published online July 19 in the *Journal of Clinical Pharmacology*.

Shih-Han Hung, M.D., from Taipei Medical University Hospital in Taiwan, and colleagues conducted a case-control study to examine the correlation of [thyroid cancer](#) with primary [hypothyroidism](#) using data from the Taiwan Longitudinal Health Insurance Database 2005. Data were included for 1,285 patients with cancer and 3,855 sex- and age-matched controls.

The researchers identified a prior hypothyroidism diagnosis in 2.88 percent of cases and 0.86 percent of controls. Compared with controls, cases were more likely to have been previously diagnosed with hypothyroidism (adjusted odds ratio, 3.01). There was a significant association for thyroid cancer with hypothyroidism patients who were regular thyroxine users (adjusted odds ratio, 8.68). For hypothyroidism patients who were irregular thyroxine users, there was no significant association with thyroid [cancer](#).

"Additional studies are needed to further evaluate and validate these findings," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Thyroid cancer tied to regular thyroxine use in hypothyroidism (2017, July 31)
retrieved 1 May 2024 from
<https://medicalxpress.com/news/2017-07-thyroid-cancer-tied-regular-thyroxine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.