

Study shows adolescent depression increases risk for violence

August 1 2017



Credit: George Hodan/Public Domain

Adolescent depression increases the risk of violence, suggests a study published in the August 2017 issue of the *Journal of the American Academy of Child and Adolescent Psychiatry (JAACAP)*.

The study examined the longitudinal association between [depression](#) and subsequent [violence](#) from three representative samples in the

Netherlands, United Kingdom and Finland. Researchers for these three cohorts used complementary measures of depression, including self-report and clinical diagnoses, and different measurements of violence including informant reports of violence and official convictions for [violent crimes](#).

The research team, led by Professor Seena Fazel, from the Forensic Psychiatry Group at the University of Oxford, United Kingdom, found modest increases in risk of violence in depression. In absolute terms, for instance, in the Finnish [sample](#), 7.1% of individuals with depression were convicted of one or more violent crimes, compared with 3.6% in the general population without depression.

In relative terms, across samples and measurements, the study shows a consistent pattern of increased relative risk of later violence. In the Dutch and UK samples, an increase in depressive symptoms was associated with a significant elevated risk of later violence. In the Finnish sample, the odds of violence in individuals with a diagnosis of depression were increased two-fold, compared to those without depression. These findings highlight the need for active and early treatment of depression in adolescents and [young people](#). The mechanisms behind this link need further investigation, and may involve increased impulsivity, hostility and poor self-regulation.

"We know that high rates of depression have been reported among adolescents in juvenile detention and correctional facilities (e.g., 11% in boys and 29% in girls)," said Dr. Rongqin Yu, lead researcher at the Forensic Psychiatry Group at the University of Oxford, United Kingdom. "However, the longitudinal link between depression and violence was unclear. Our longitudinal design allowed us to take into account previous violence, enabling us to test whether [adolescent depression](#) is associated with changes in violence over time. We found a consistent pattern of increased risk of later violence across samples. Both

depression and violence are prevalent in adolescents and young adults; our findings indicate the importance of early detection and treatment of depression."

Professor Fazel said: "This research is important for two main reasons. First, it adds to the evidence of the many potential harms of untreated depression in young people. Second, it suggests that closer liaison between criminal justice and mental health might prevent violence in high-risk individuals."

More information: Rongqin Yu et al. Depression and Violence in Adolescence and Young Adults: Findings From Three Longitudinal Cohorts, *Journal of the American Academy of Child & Adolescent Psychiatry* (2017). [DOI: 10.1016/j.jaac.2017.05.016](https://doi.org/10.1016/j.jaac.2017.05.016)

Provided by Elsevier

Citation: Study shows adolescent depression increases risk for violence (2017, August 1) retrieved 4 May 2024 from <https://medicalxpress.com/news/2017-08-adolescent-depression-violence.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
