

Age-related hearing loss and communication breakdown in the clinical setting

August 24 2017

It was not uncommon for older adults to report mishearing a physician or nurse in a primary care or hospital setting, according to a study published by *JAMA Otolaryngology-Head & Neck Surgery*.

The prevalence of [medical errors](#) is higher among older patients. Failures in clinical communication are considered to be the leading cause of medical errors. A previous study reported that improved communication between the medical teams and families could have prevented 36 percent of medical errors. Colm M. P. O'Tuathaigh, B.A., Ph.D., of University College Cork, Cork, Ireland and colleagues conducted an analysis of interview data collected in 100 adults 60 years and older to examine [communication breakdown](#) in hospital and [primary care](#) settings among adults reporting hearing loss.

Of these adults, 57 reported some degree of hearing loss; 26 used a hearing aid device. Of the 100 adults, 43 reported having misheard a physician, nurse or both in a primary care or [hospital setting](#). When asked to elaborate on the context of mishearing in a clinical setting, the scenarios included (in descending order of citation frequency): general mishearing, consultation content, physician-patient or nurse-patient communication breakdown, hospital setting and use of language.

"This qualitative analysis confirms that age-related [hearing loss](#) has a negative effect on clinical communication across both hospital and primary care clinical settings," the authors write. "We recommend that content-related and setting-related factors identified as barriers to

communication in adults with hearing impairment be incorporated within a patient-centered approach to clinical communication with this patient population."

More information: *JAMA Otolaryngology-Head & Neck Surgery*, [jamanetwork.com/journals/jamaoto ... 01/jamaoto.2017.1248](https://jamanetwork.com/journals/jamaoto/01/jamaoto.2017.1248)

Provided by The JAMA Network Journals

Citation: Age-related hearing loss and communication breakdown in the clinical setting (2017, August 24) retrieved 4 May 2024 from <https://medicalxpress.com/news/2017-08-age-related-loss-breakdown-clinical.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--