

Antidepressant use increases risk of head injuries among persons with Alzheimer's disease

August 9 2017

Antidepressant use is associated with an increased risk of head injuries and traumatic brain injuries among persons with Alzheimer's disease, according to a new study from the University of Eastern Finland.

Antidepressant use has previously been linked with an increased risk of falls and hip fractures, but the risk of head injuries has not been studied before. The results were published in *Alzheimer's Research & Therapy*.

Antidepressant use was associated with a higher risk of [head injuries](#) especially at the beginning of use – during the first 30 days –, but the risk persisted even longer, up to two years. The association was also confirmed in a study design comparing time periods within the same person, thus eliminating selective factors. The association with traumatic brain injuries was not as clear as for head injuries, which may be due to a smaller number of these events in the study population. The use of other psychotropic drugs did not explain the observed associations.

Head injuries are more common among older people than younger ones, and they are usually caused by falling. As antidepressant use has previously been associated with an increased risk of falling, the researchers were not surprised that the use of antidepressants also increased the risk of head injuries.

"However, our findings give cause for concern because persons with Alzheimer's disease frequently use antidepressants, which have been

considered a safer alternative to, for example, benzodiazepines," says Senior Researcher Heidi Taipale from the University of Eastern Finland.

"Our study population consisted of persons diagnosed with Alzheimer's disease, but it is likely that the [risk](#) is similar also in other older persons without Alzheimer's disease. This is something we will be studying in the future."

The study constitutes part of the nationwide register-based MEDALZ study, which includes all community-dwelling persons diagnosed with Alzheimer's disease in Finland during 2005-2011. The study included 10,910 antidepressant users and 21,820 nonusers, all of whom had Alzheimer's disease.

More information: Heidi Taipale et al. Risk of head and traumatic brain injuries associated with antidepressant use among community-dwelling persons with Alzheimer's disease: a nationwide matched cohort study, *Alzheimer's Research & Therapy* (2017). [DOI: 10.1186/s13195-017-0285-3](https://doi.org/10.1186/s13195-017-0285-3)

Provided by University of Eastern Finland

Citation: Antidepressant use increases risk of head injuries among persons with Alzheimer's disease (2017, August 9) retrieved 17 April 2024 from <https://medicalxpress.com/news/2017-08-antidepressant-injuries-persons-alzheimer-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--