


Cancer detection with sugar molecules

August 14 2017



Like a spaceship, the complex sugar molecule (coloured) lands exactly on the tumor protein galectin-1, which here looks like a meteorite and is shown in black



and white. Credit: Workgroup Seibel, VCH-Wiley

Galectins are a family of proteins that have become a promising source of cancer research in recent years. A representative thereof is galectin-1. It sits on the surface of all human cells; on tumor cells, however, it occurs in enormous quantities. This makes it an interesting target for diagnostics and therapy.

"Among other things, it is known that galectin-1 hides the <u>tumor cells</u> from the immune system," explains Professor Jürgen Seibel of the Institute of Organic Chemistry at the Julius-Maximilians-Universität (JMU) Würzburg in Bavaria, Germany. Recent studies have shown that when Galectin-1 is blocked, the immune system can recognize the tumor and attack it with T cells.

Sugar molecule with docking station

No wonder, therefore, that galectin-1 has become a major focus of research. Seibel and his colleague Dr. Clemens Grimm is interested in a very specific section of this protein, the so-called carbohydrate recognition domain. They have now designed a complex sugar molecule that fits perfectly into this domain, as the scientists report in journal *ChemBioChem*.

"We have equipped the sugar molecule with a docking site, for example, to connect it with a fluorescent dye or an drug," says Seibel. In addition, the scientists have described the binding of their molecule to galectin-1 with high-resolution X-ray structure analyzes.

"Our findings can serve the development of high-affinity ligands of the protein Galectin-1 and thus of new drugs," said Clemens Grimm.



Quick test for Galectin-1 in progress

Now the JMU scientists are working on a rapid test for the detection of galectin-1. It is designed to enable early detection of tumors such as neuroblastoma. For the future, Seibel's team would like to expand the <u>sugar molecules</u> into a kind of shuttle system that allows pharmaceutical agents to be transported directly to the tumors.

More information: Nadja Bertleff-Zieschang et al, Exploring the Structural Space of the Galectin-1-Ligand Interaction, *ChemBioChem* (2017). DOI: 10.1002/cbic.201700251

Provided by University of Würzburg

Citation: Cancer detection with sugar molecules (2017, August 14) retrieved 18 April 2024 from https://medicalxpress.com/news/2017-08-cancer-sugar-molecules.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.