

# A faster, less costly way to process walnuts

August 21 2017, by Sandra Avant

---



ARS engineer Zhongli Pan (right) and post docs Ragab Khir (center) and Chandrasekar Venkitasamy use an infrared thermometer to check walnut temperatures. Credit: Mathew Reiff

As part of a healthful diet, walnuts provide protein, antioxidants, essential vitamins, and minerals. Walnuts are also a good source of omega-3 fatty acids, which are associated with overall good health.

However, before walnuts reach your grocery store, they must be harvested, washed, dehulled (hulls removed from shells), and dried. These processes use a lot of energy.

Agricultural Research Service (ARS) scientists in Albany, California, have developed a method to improve [walnut](#) processing, particularly drying, to reduce energy and obtain high-quality walnut products.

California produces 99 percent of U.S. walnuts—about 570,000 tons per year, says engineer Zhongli Pan, who is in ARS's Healthy Processed Foods Research Unit in Albany.

Drying walnuts with hot air takes more than 24 hours and uses a significant amount of natural gas and electricity. "The current hot-air drying method is very slow and not efficient," Pan says. "We knew that infrared had a higher energy transfer rate and could dry walnuts faster, giving us an opportunity to improve the process."

Infrared is a type of electromagnetic radiation, as are radio waves, ultraviolet radiation, x rays, and microwaves. Infrared is a form of light that we cannot see—but we can feel its heat on our skin.



ARS engineer Zhongli Pan (right), ARS post doc Chandrasekar Venkitasamy (left) and Justin McCurdy of Wizard Manufacturing (center) check walnuts drying with infrared heat. Credit: Mathew Reiff

Pan and his research team discovered that up to 25 percent of the energy used to dry walnuts could be saved by pre-drying them with infrared heat, followed by regular hot-air drying.

After washing, walnuts have a broad range of [moisture content](#), says Pan, but currently they are hot-air dried for the same amount of time. This can cause underdrying or overdrying.

"If the [moisture](#) content is too high, it will cause mold," Pan says, "so facilities tend to overdry walnuts to make sure they are safe to



consume." Overdrying wastes a lot of energy and reduces walnut quality.

The new infrared-drying process quickly removes moisture from the surface of walnuts, which shortens total drying time by 35 percent. It reduces overdrying and underdrying and allows the walnuts inside the shells to maintain their color and exceptional polyunsaturated fat content.

Says Pan, "The good thing is that we reduced the surface moisture content of walnuts quickly," which prevents the moisture from moving into the walnut. "It not only saves [energy](#), but it's a safer and better way to dry walnuts." This new technology may also be feasible for other types of nuts, such as pistachios.



Walnuts. Credit: ARS

Provided by Agricultural Research Service

Citation: A faster, less costly way to process walnuts (2017, August 21) retrieved 26 June 2024 from <https://medicalxpress.com/news/2017-08-faster-costly-walnuts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.