

Higher coffee consumption associated with lower risk of death

August 28 2017



Credit: George Hodan/public domain

Higher coffee consumption is associated with a lower risk of death, according to research presented today at ESC Congress. The observational study in nearly 20 000 participants suggests that coffee can be part of a healthy diet in healthy people.

"Coffee is one of the most widely consumed beverages around the

world," said Dr Adela Navarro, a cardiologist at Hospital de Navarra, Pamplona, Spain. "Previous studies have suggested that drinking coffee might be inversely associated with all-cause mortality but this has not been investigated in a Mediterranean country."

The purpose of this study was to examine the association between [coffee consumption](#) and the risk of mortality in a middle-aged Mediterranean cohort. The study was conducted within the framework of the Seguimiento Universidad de Navarra (SUN) Project, a long-term [prospective cohort study](#) in more than 22 500 Spanish university graduates which started in 1999.

This analysis included 19 896 participants of the SUN Project, whose average age at enrolment was 37.7 years old. On entering the study, participants completed a previously validated semi-quantitative food frequency questionnaire to collect information on coffee consumption, lifestyle and sociodemographic characteristics, anthropometric measurements, and previous health conditions.

Patients were followed-up for an average of ten years. Information on mortality was obtained from study participants and their families, postal authorities, and the National Death Index. Cox regression models were used to estimate hazard ratios (HR) and 95% confidence intervals (CI) for incident mortality according to baseline total coffee consumption adjusted for potential confounders.

During the ten year period, 337 participants died. The researchers found that participants who consumed at least four cups of coffee per day had a 64% lower risk of all-cause mortality than those who never or almost never consumed coffee (adjusted HR, 0.36; 95% CI, 0.19-0.70). There was a 22% lower risk of all-cause mortality for each two additional cups of coffee per day (adjusted HR, 0.78; 95% CI, 0.66-0.92).

The researchers examined whether sex, age or adherence to the Mediterranean diet had any influence on the association between baseline coffee consumption and mortality. They observed a significant interaction between coffee consumption and age (p for interaction=0.0016). In those who were at least 45 years old, drinking two additional cups of coffee per day was associated with a 30% lower risk of mortality during follow-up (adjusted HR, 0.70; 95% CI, 0.58-0.85). The association was not significant among younger participants.

Dr Navarro said: "In the SUN project we found an inverse association between drinking coffee and the risk of all-cause mortality, particularly in people aged 45 years and above. This may be due to a stronger protective association among older participants."

She concluded: "Our findings suggest that drinking four cups of [coffee](#) each day can be part of a [healthy diet](#) in healthy people."

Provided by European Society of Cardiology

Citation: Higher coffee consumption associated with lower risk of death (2017, August 28) retrieved 5 May 2024 from

<https://medicalxpress.com/news/2017-08-higher-coffee-consumption-death.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--