

# Lack of REM sleep may lead to higher risk for dementia

August 23 2017

---



Credit: CC0 Public Domain

Spending less time in REM (rapid eye movement) sleep and taking longer to enter REM sleep are separately associated with a higher risk of developing dementia.

These findings appear in the journal *Neurology*, the medical journal of the American Academy of Neurology," and were presented in July at the Alzheimer's Association International Conference in London.

It is common for people with dementia to experience sleep disturbance. However, it is unclear if sleep disturbance occurs as a consequence of dementia or if disturbed sleep is associated with the risk of dementia in the future.

Using data from the Framingham Heart Study (FHS), researchers from Boston University School of Medicine (BUSM) studied 321 participants over the age of 60 who underwent an overnight sleep study between 1995 and 1998. They were then followed for an average of 12 years to determine their risk of developing dementia. Upon follow-up, the researchers found that each percentage reduction in REM sleep was associated with a nine percent increase in the risk of all-cause dementia and an eight percent increase in the risk of Alzheimer's disease dementia.

"Different stages of sleep may differentially affect key features of Alzheimer's disease. Our findings implicate REM sleep mechanisms as predictors of dementia," explained Matthew Pase, PhD, fellow in the department of [neurology](#) at BUSM and investigator at the FHS who presented the findings.

Earlier this year, Pase and his colleagues found that people who consistently sleep more than nine hours each night had double the risk of developing dementia in 10 years as compared to participants who slept for nine hours or less. Those findings were reported in the journal *Neurology*.

According to the researchers future research is needed to determine whether REM sleep helps protect the brain from dementia or is sensitive

to early brain changes that accompany [dementia](#).

Provided by Boston University School of Medicine

Citation: Lack of REM sleep may lead to higher risk for dementia (2017, August 23) retrieved 23 April 2024 from <https://medicalxpress.com/news/2017-08-lack-rem-higher-dementia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.