

New type of MRI scan developed to predict stroke risk

August 24 2017



New scan developed to predict stroke risk. Credit: Martin Phelps

Researchers at the University of Oxford have developed a new type of MRI scan to predict the risk of having a stroke, thanks to funding from the British Heart Foundation (BHF).

The non-invasive technique, described in a paper published in the journal *JACC: Cardiovascular Imaging*, produces a quantitative result that can accurately indicate whether plaques in the [carotid](#) arteries – those that supply the brain with blood - are rich in cholesterol, and

therefore more likely to cause a stroke.

The rupture of [fatty plaques](#) can block the arteries and cause potentially debilitating and life-threatening strokes as the brain is starved of oxygen. There are more than 100,000 strokes in the UK each year, around a quarter of which are caused by carotid plaques.

Latest figures show that stroke now costs the UK healthcare system an estimated £2 billion every year. It is estimated that stroke costs the UK as much as £9 billion a year as a society.

At present, the risk of stroke is measured by the size of the [plaque](#) in the [carotid artery](#). If the plaque is deemed to be too big, people are treated surgically to remove it. However, this method can miss fatty plaques that are not big, but have a high risk of rupturing.

The new MRI technique was developed to differentiate between the risky plaques that contain a lot of cholesterol, and those that are more stable.

In the study, the researchers used the new MRI scan to measure the amount of cholesterol in the carotid plaques of 26 patients scheduled for surgery. After the plaques were surgically removed, the team looked at the actual cholesterol content in each plaque and found that the new technique was accurate and the more cholesterol they detected within the plaque, the greater the risk.

The work was a collaboration between researchers at the University of Oxford and surgeons working within the John Radcliffe Hospital and was supported by the BHF Centre of Research Excellence in Oxford and the NIHR Oxford Biomedical Research Centre.

One of the authors, Dr Luca Biasioli, from the University of Oxford,

said: 'Stroke is a leading cause of disability and the third biggest killer in the UK.

When someone goes to hospital having suffered a minor stroke, it's vital that doctors know whether the patient might be at risk of a further stroke, which could be fatal.

'Being able to quantify [cholesterol](#) in carotid plaques is a really exciting prospect, as this new MRI technique could help doctors to identify patients at higher risk of stroke and make more informed decisions on their treatments.'

Professor Sir Nilesh Samani, Medical Director at the BHF, which part-funded the study, said: 'When someone suffers a mini-stroke, they often go on to suffer a more serious, or even deadly, stroke in the hours, days or weeks that follow. This exciting research opens up the possibility that in the future we may be able to more accurately identify people with carotid plaques that are likely to rupture and cause a [stroke](#).

'These patients can then be treated earlier, for example with surgery to remove the plaque, while others might be spared surgery altogether. More research is now necessary before this advance can come into routine clinical practice. However, if successful this technique has the promise to save lives.'

The full paper can be read in the journal *JACC: Cardiovascular Imaging*.

More information: Joshua T. Chai et al. Quantification of Lipid-Rich Core in Carotid Atherosclerosis Using Magnetic Resonance T 2 Mapping, *JACC: Cardiovascular Imaging* (2016). [DOI: 10.1016/j.jcmg.2016.06.013](https://doi.org/10.1016/j.jcmg.2016.06.013)

Provided by University of Oxford

Citation: New type of MRI scan developed to predict stroke risk (2017, August 24) retrieved 28 April 2024 from <https://medicalxpress.com/news/2017-08-mri-scan.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.