

Opioids overused in migraine treatment, regardless of race, study finds

August 17 2017



African-Americans are more likely to experience debilitating migraine headaches than whites, but a new study probing the issue found no evidence of racial disparities in treatment practices. Credit: University of Michigan

African-Americans are more likely to experience debilitating migraine headaches than whites, but a new study probing the issue found no evidence of racial disparities in treatment practices.

Instead, researchers report a different finding that affects everyone: opioid overuse.

"Our hypothesis was not proven," says first author Larry Charleston IV, M.D., M.Sc., an assistant professor of neurology at the University of Michigan Medical School. "However, we found people were getting prescribed opioids as much as they were getting prescribed medications that are much better for migraines. This is an opportunity for clinicians to improve prescribing practices when treating [migraine](#) patients."

Charleston, along with author James Burke, M.D., also an assistant professor of neurology at U-M, studied results from 2,860 migraine visits in the U.S. between 2006 and 2013. Seventy-six percent of patients were non-Hispanic whites, while 10 percent each were African-American or Hispanic.

The authors reported the research in *Cephalalgia*, the journal of the International Headache Society.

Evidence-based medications underused

African-Americans were 4 percent less likely to receive all high-quality abortive agents and Hispanics were 5 percent less likely to receive all high-quality prophylactic agents compared with non-Hispanic whites, yet neither difference was statistically significant. Charleston says evidence-based medications are being used about as often as opioids.

In all groups, 15 to 20 percent of patients received a Level A abortive, which is an evidence-based medication intended for acute migraine treatment, such as a triptan or dihydroergotamine. (Specifically, 15.3 percent of African-Americans, 19.4 percent of non-Hispanic whites and 17.7 percent of Hispanics received one.)

Overall, 38.9 percent of patients were not prescribed any abortive medication.

"This shows an underuse of medications with high-quality evidence," Charleston says. "Even for patients being prescribed an abortive medication, we found 27 percent of them were given at least one low-quality abortive medication. Better options do exist."

And more than 40 percent of patients studied received no prophylactic treatment, which are preventive medications (41.3 percent of African-Americans, 40.8 percent of non-Hispanic whites and 41.2 percent of Hispanics).

Across the board, 15.2 percent of patients studied received a prescription for an opioid, with no significant difference among African-American, Hispanic and non-Hispanic white patients. In addition to the evidence that shows other medications are simply more effective for treating migraine headaches, opioids carry additional problems including association with [medication](#) overuse [headache](#) and comorbidities such as depression and anxiety.

Racial disparities in migraines

Migraine headaches are a huge problem, costing the country billions of dollars and affecting quality of life for those who have them. In fact, the American Academy of Neurology reports that more than a quarter of people with migraine headaches have missed at least one day of work in the past three months.

"We already knew the migraine burden is increased in African-American [patients](#)," says Charleston, a member of U-M's Institute for Healthcare Policy and Innovation. "African-Americans experience more frequent and severe migraine headaches than non-Hispanic whites, yet

they respond to treatment just as well."

Charleston says the research was designed to investigate these disparities in migraine occurrence and severity. But because this study was not able to determine the reasons for [racial disparities](#) in [migraine headaches](#), he says future research is needed.

More information: Larry Charleston IV et al, Do racial/ethnic disparities exist in recommended migraine treatments in US ambulatory care?, *Cephalalgia* (2017). [DOI: 10.1177/0333102417716933](#)

Provided by University of Michigan

Citation: Opioids overused in migraine treatment, regardless of race, study finds (2017, August 17) retrieved 21 July 2024 from <https://medicalxpress.com/news/2017-08-opioids-overused-migraine-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--