


Poor sleep is associated with ischemic heart disease and stroke

August 29 2017



Proportion of poor sleep in ischaemic heart disease (IHD), stroke, and non-cardiovascular disease (CVD) populations. Poor sleep was defined as a global Pittsburgh Sleep Quality Index score of 6 or more. Credit: European Society of Cardiology

Poor sleep is associated with ischaemic heart disease and stroke, according to research presented today at ESC Congress. The observational study in nearly 13 000 people revealed different patterns of sleep disturbance between the two conditions, with ischaemic heart disease being linked to shorter sleep and brief moments of waking up.


"Poor sleep is associated with cardiovascular diseases such as [ischaemic heart disease](#) and [stroke](#) but the kind of [sleep disturbances](#) that are most

risky is not well documented," said lead researcher Dr Nobuo Sasaki, of the Hiroshima Atomic Bomb Casualty Council, Japan. "'Poor sleep' includes too short or too long sleep, difficulty falling asleep, and difficulty maintaining sleep."

This study investigated the association between sleep disturbances and [cardiovascular disease](#). It also aimed to clarify possible differences in sleep disturbances between ischaemic heart disease and stroke.

The study included 12 876 residents of Hiroshima, Japan (6 762 men and 6 114 women, average age 68 years) who were registered for an annual health check. Of those, 773 patients had a history of ischaemic heart disease (myocardial infarction and/or angina), 560 patients had a history of stroke (intracranial haemorrhagic and/or cerebral infarction), and 11 543 had no cardiovascular disease. Patients with both ischaemic heart disease and stroke, or another type of cardiovascular disease, were excluded from the study.

Sleep habits were assessed with the Pittsburgh Sleep Quality Index (PSQI), a 19-item self-reporting questionnaire which yields seven component scores. C1 assesses subjective [poor sleep](#) quality, C2 long sleep latency, C3 short sleep duration, C4 low sleep efficiency, C5 difficulty in maintaining sleep, C6 use of sleeping pills, and C7 daytime dysfunction. Each component is ranked 0, 1, 2, or 3, with a score ≥ 2 defining sleep disturbance (except C6 score ≥ 1).



Proportions of subjective sleep quality (C1), long sleep latency (C2), short sleep duration (C3), low sleep efficiency (C4), difficulty maintaining sleep (C5), use of sleeping pills (C6), and daytime dysfunction (C7) in ischaemic heart disease (IHD), stroke, and non-cardiovascular disease (CVD) populations. Each type of disturbed sleep was defined as a score ≥ 2 (except C6 score ≥ 1). Credit: European Society of Cardiology

A sum of the seven scores was used to calculate the global PSQI score which ranged from 0 to 21. Higher scores indicated poorer sleep quality, and 'poor sleep' was defined as a global PSQI score ≥ 6 .

Rates of 'poor sleep' and component sleep disturbances are shown in figures A and B. Poor sleep occurred in 52%, 48%, and 37% of patients with ischaemic heart disease, stroke, and no cardiovascular disease, respectively.

After adjusting for confounding factors (table) 'poor sleep' was significantly associated with ischaemic heart disease (odds ratio [OR], 1.71; p

Citation: Poor sleep is associated with ischemic heart disease and stroke (2017, August 29) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2017-08-poor-ischemic-heart-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.