

Reconstruction method in gastric CA surgery affects bone density

August 4 2017



(HealthDay)—The reconstruction method may affect postoperative bone

mineral density (BMD) loss in gastric cancer, according to a study published online July 31 in the *Journal of Gastroenterology and Hepatology*.

Taisuke Imamura, M.D., from the Kyoto Prefectural University of Medicine in Japan, and colleagues assessed 148 consecutive patients who underwent distal gastrectomy with Billroth-I (B-I) or Roux-en-Y (R-Y) reconstruction for stage I [gastric cancer](#) (2008 to 2012). BMD data were retrospectively assessed using computed tomography attenuation values of the first lumbar vertebra after surgery.

In multivariate analysis, the researchers found that R-Y reconstruction was an [independent risk factor](#) for BMD loss after distal gastrectomy (P

"B-I [reconstruction](#) might be a preferable method for preventing BMD loss after gastrectomy in gastric [cancer](#) patients," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Reconstruction method in gastric CA surgery affects bone density (2017, August 4) retrieved 5 May 2024 from <https://medicalxpress.com/news/2017-08-reconstruction-method-gastric-ca-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--