


Scientists develop infection model for tickborne flaviviruses

August 22 2017



Langkat virus infection (bright green) in the tick midgut (black) is shown at six days after infection in this fluorescence image. Credit: NIAID

National Institutes of Health (NIH) scientists have filled a research gap by developing a laboratory model to study ticks that transmit flaviviruses, such as Powassan virus. Powassan virus was implicated in the death of a New York man earlier this year. The unusual model involves culturing organs taken from *Ixodes scapularis* ticks and then infecting those organ cultures with flaviviruses, according to researchers at Rocky Mountain Laboratories, part of NIH's National Institute of Allergy and Infectious Diseases (NIAID). The researchers say the culture model will greatly increase knowledge about how flaviviruses infect ticks and could become a tool to evaluate medical countermeasures against tick-borne viruses.

Flaviviruses are the cause of diseases spread by mosquitoes (e.g., [dengue fever](#) and West Nile fever) and by ticks (e.g., Powassan virus disease and tick-borne encephalitis). Powassan virus and the closely related deer tick virus are the only flaviviruses known to be spread by ticks in North America. In the last 10 years, about 75 U.S. cases of Powassan virus infection have been reported, according to the Centers for Disease Control and Prevention. Powassan virus infection can result in fever, headache, vomiting, weakness, confusion, seizures, memory loss, and death. No licensed treatments or vaccines are available for Powassan virus disease.

The NIAID scientists developed their model by dissecting three tick organs—the midgut, salivary glands and nervous tissue—and then culturing [flaviviruses](#) in those organs, evaluating their viability over several days. They found that Powassan virus and the related Langat virus could infect and spread in salivary glands and midgut. Langat virus is found typically in Southeast Asia and is an ideal model [virus](#) for study because it causes only rare, mild infections in people.

More information: J Grabowski et al. Flavivirus infection of *Ixodes scapularis* (black-legged tick) ex vivo organotypic cultures and

application for control. *mBio*. DOI: [10.1128/mBio.01255-17](https://doi.org/10.1128/mBio.01255-17)

Provided by NIH/National Institute of Allergy and Infectious Diseases

Citation: Scientists develop infection model for tickborne flaviviruses (2017, August 22)
retrieved 26 April 2024 from

<https://medicalxpress.com/news/2017-08-scientists-infection-tickborne-flaviviruses.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.