

Sonography may aid hyaluronic acid injections for knee arthritis

August 14 2017



(HealthDay)—Sonography-guided hyaluronic acid injection may yield

better results than surface anatomy-guided injection in patients with knee osteoarthritis, according to a study published online Aug. 9 in the *International Journal of Rheumatic Diseases*.

Nahid Kianmehr, M.D., from Iran University in Tehran, and colleagues compared the effect of sonographic-guided versus blind knee injection of hyaluronic acid among 61 patients with primary [knee osteoarthritis](#).

The researchers found that all scores in the Western Ontario and McMaster Universities Osteoarthritis Index pain and function subscales changed significantly in both short- and long-term follow-up. However, in those who had received intra-articular injection under the guidance of sonography, the mean differences at six and 12 weeks were significantly higher for both the pain and function subscales. At six weeks, the mean in differences of the 10-cm visual analogue scale (both at rest and after 50-feet walk) was statistically significant in both groups, but did not remain significant after 12 weeks.

"Application of sonography might improve the response of patients to intra-articular injection of [hyaluronic acid](#), at least in certain clinical indices," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Sonography may aid hyaluronic acid injections for knee arthritis (2017, August 14) retrieved 10 April 2024 from <https://medicalxpress.com/news/2017-08-sonography-aid-hyaluronic-acid-knee.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.