

Which vaccinations should I get as an adult?

August 29 2017, by C. Raina Macintyre And Rob Menzies



Credit: AI-generated image ([disclaimer](#))

Before vaccines were developed, infectious diseases such as diphtheria, tetanus and meningitis were the [leading cause of death](#) and illness in the world. Vaccines are one of the [greatest public health achievements in history](#), having drastically reduced deaths and illness from infectious causes.

There is a large [gap between vaccination rates](#) for funded [vaccines for](#)

[adults in Australia](#) and those for infants. More than 93% of infants are vaccinated in Australia, while in adults the rates are between 53-75%. Much more needs to be done to prevent infections in adults, particularly those at risk.

If you are an adult in Australia, the kinds of vaccines you need to get will depend on several factors, including whether you missed out on childhood vaccines, if you are Aboriginal or Torres Strait Islander, your occupation, how old you are and whether you intend to go travelling.

For those born in Australia

Children up to four years and aged 10-15 receive vaccines under the [National Immunisation Schedule](#). These are for hepatitis B, whooping cough, diphtheria, tetanus, measles, mumps, rubella, polio, haemophilus influenzae B, rotavirus, pneumococcal and meningococcal disease, chickenpox and the human papillomavirus (HPV).

Immunity following vaccination varies depending on the vaccine. For example, the measles vaccine protects for a long duration, possibly a lifetime, whereas immunity wanes for pertussis (whooping cough). Boosters are given for many vaccines to improve immunity.

Measles, mumps, rubella, chickenpox, diphtheria and tetanus

People born in Australia before 1966 likely have [natural immunity to measles](#) as the viruses were circulating widely prior to the vaccination program. People born after 1965 should have received two doses of a [measles vaccine](#). Those who haven't, or aren't sure, can safely receive a vaccine to avoid infection and prevent transmission to babies too young to be vaccinated.

Measles vaccine can be given as MMR (measles-mumps-rubella) or MMRV, which includes varicella (chickenpox). The [varicella vaccine](#) on its own (not combined in MMRV) is advised for people aged 14 and over who have not had chickenpox, especially women of childbearing age.

Booster doses of diphtheria, tetanus and whooping cough vaccines, are available free at age 10-15, and recommended at 50 years old and also at 65 years and over if not received in the previous ten years. Anyone unsure of their tetanus vaccination status who sustains a tetanus-prone wound (generally a deep puncture or wound) should get vaccinated. While tetanus is rare in Australia, most cases we see are in older adults.




Whooping cough

Pregnant women are recommended to get the [diphtheria-tetanus-acellular pertussis](#) vaccine in the third trimester to protect the vulnerable infant after it is born, and [influenza vaccine](#) at any stage of the pregnancy (see below under influenza).

Pertussis (whooping cough) is a contagious respiratory infection dangerous for babies. One in [every 200 babies](#) who contract [whooping cough](#) will die.

It is particularly important for women from 28 weeks gestation to ensure they are vaccinated, as well as the partners of these women and anyone else who is taking care of a child younger than six months old. Deaths from pertussis are also documented in elderly Australians.

Vaccinations for adults: the easy wins

Disease	What is it?	Who should get it?
Measles	A contagious viral illness that causes skin rash and fever, which can lead to encephalitis and death. 	Anyone who hasn't had two doses, or isn't sure if they have, should get the measles-mumps-rubella (MMR).
Mumps	A viral illness that causes fever and swollen salivary glands. 	
German measles (Rubella)	A viral illness that causes a skin rash and joint pain. Can cause death or birth defects in an unborn baby. 	
Diphtheria	A bacterial disease that causes severe inflammation of the nose, throat and windpipe (trachea), which can lead to suffocation.	Adolescents aged 10 to 15, adults at 50 and those over 65 who have not had a booster in more than 10 years should get the diphtheria-tetanus-acellular pertussis (DTPa-containing) booster vaccine. Any deep wound could be prone to tetanus, so check with a doctor to see if you need a precautionary booster for tetanus. Women over 28 months pregnant (and their partners). Parents or guardians of children under 6 months old should also make sure they're up-to-date.
Tetanus	A bacterial disease that causes muscle spasms and breathing problems, generally caused by wound contamination from soil, dust or manure. Can be fatal.	
Whooping cough (Pertussis)	A very contagious respiratory infection with a characteristic cough followed by a whooping sound during inhalation. Often fatal for infants.	



In July 2017, the government announced free catch-up vaccinations for all newly arrived refugees. This covers any childhood vaccine on the National Immunisation Schedule which has been missed. Credit: Information sourced from betterhealth.vic.gov.au and [healthdirect.gov.au/The Conversation](http://healthdirect.gov.au/TheConversation), CC BY-ND

Pneumococcal disease and influenza

The [pneumococcal vaccine](#) is funded for everyone aged 65 and over, and [recommended for anyone](#) under 65 with risk factors such as chronic lung disease.

Anyone from the age of six months can get the flu (influenza) vaccine. The vaccine can be given to any adult who requests it, but is only funded if they fall into defined risk groups such as pregnant women, Indigenous Australians, people aged 65 and over, or those with a medical condition such as chronic lung, cardiac or kidney disease.

Flu vaccine is matched every year to the anticipated circulating flu viruses and is quite effective. The vaccine covers four strains of influenza. Pregnant women are at increased risk of the flu and recommended for influenza vaccine any time during pregnancy.

Health workers, childcare workers and aged-care workers are a priority for vaccination because they care for sick or vulnerable people in institutions at risk of outbreaks. Influenza is the most important vaccine for these occupational groups, and some organisations provide free staff vaccinations. Otherwise, you can ask your doctor for a vaccination.

Any person whose immune system is [weakened through medication](#) or illness (such as HIV) is at increased risk of infections. However, live

viral or bacterial vaccines must not be given to immunosuppressed people. They must seek medical advice on which vaccines can be safely given.

Hepatitis

Australian-born children receive four shots of the hepatitis B vaccine, but some adults are advised to get vaccinations for hepatitis A or B.

Those recommended to receive the [hepatitis A vaccine](#) are: travellers to hepatitis A endemic areas; people whose jobs put them at risk of acquiring hepatitis A including childcare workers and plumbers; men who have sex with men; injecting drug users; people with developmental disabilities; those with chronic liver disease, liver organ transplant recipients or those chronically infected with hepatitis B or hepatitis C.

Those recommended to get the hepatitis B vaccine are: people who live in a household with someone infected with hepatitis B; those having sexual contact with someone infected with hepatitis B; sex workers; men who have sex with men; injecting drug users; migrants from hepatitis B endemic countries; healthcare workers; Aboriginal and Torres Strait Islanders; and some others at high risk at their workplace or due to a medical condition.

Human papillomavirus

The [human papillomavirus \(HPV\) vaccine](#) protects against cervical, anal, head and neck cancers, as well as some others. It is available for boys and girls and delivered in high school, usually in year seven. There is benefit for older girls and women to be vaccinated, at least up to their mid-to-late 20s.

The elderly

With ageing comes a progressive decline in the immune system and a corresponding increase in risk of infections. Vaccination is the [low-hanging fruit](#) for healthy ageing. The elderly are advised to receive the influenza, pneumococcal and shingles vaccines.

Influenza and pneumonia are [major preventable causes](#) of illness and death in older people. The flu causes deaths in children and the elderly during severe seasons.

Vaccinations for adults: individual diseases

Disease	What is it?	Who should get it?
Pneumococcal disease	Range of infections in various body parts. Illnesses range from mild infections, such as ear infection, to pneumonia and life-threatening infections of the bloodstream and central nervous system, such as meningitis.	People over 65 years old and people with impaired immune systems. Indigenous Australians are also at increased risk. Funded for all people aged 65 and over, and Indigenous people over the age of 50 or between 15-49 years with medical risk factors.
Human papillomavirus (HPV)	The human papillomavirus (HPV) is responsible for most cancers of the cervix, it has also been associated with cancers of the anus, head and neck, as well as some others.	All boys and girls should be vaccinated against HPV, and vaccinations are free as part of the school program. For adults, there is benefit for women to be vaccinated until the mid to late 20s.
Shingles	Caused by the varicella zoster virus, which is also responsible for chickenpox. Forms a rash that develops into very painful blisters, generally on one side of the body, face or pelvis.	Shingles can lead to debilitating and chronic pain in older people, as well as dangerous infections. A single dose of the zoster vaccine is recommended for those aged 60 and onwards, and free from 70-79 years old.
Hepatitis A	Viral disease that affects the liver, causing fever, nausea, and yellow skin and eyes. Spread through contact with food, drinks or objects contaminated by the faeces of an infected person.	Travellers to developing countries or endemic areas, childcare workers, IV drug users, sex workers, people with chronic liver disease, those already infected with hepatitis B or C and Indigenous Australians. Can be vaccinated individually or with a combined A/B vaccine.
Hepatitis B	Viral infection that causes liver inflammation, loss of appetite, fever and pain in the joints. Can lead to serious long-term complications such as cirrhosis of the liver and cancer. Spread through body fluid contact, such as sexual contact or sharing injection equipment.	People who live in a household with someone with hepatitis B, those having sexual contact with someone with hepatitis B, sex workers, men who have sex with men, IV drug users, migrants from hepatitis B endemic countries, health care workers and Indigenous Australians.
Flu (Influenza)	A highly contagious viral infection spread by contact with fluids from coughs or sneezes from an infected person. Can cause a runny and stuffy nose, sore throat, fever, cough, loss of appetite, body aches and more. Can lead to serious complications including pneumonia and death.	Every adult can get vaccinated annually, but people over 65, Indigenous Australians, pregnant women, and those with conditions like severe asthma, diabetes, and lung or heart disease are particularly at risk and vaccination is free for these groups. Many workplaces also provide free vaccination.
Chicken pox (Varicella)	Highly contagious viral disease which causes an itchy, blistering skin rash. Complications can be more severe in adults than children.	The varicella vaccine is recommended for all adults who've not previously had chicken pox.



In July 2017, the government announced free catch-up vaccinations for all newly arrived refugees. This covers any childhood vaccine on the National Immunisation Schedule which has been missed. Credit: Information sourced from betterhealth.vic.gov.au and [healthdirect.gov.au/The Conversation](http://healthdirect.gov.au/The_Conversation), CC BY-ND

The most common cause of pneumonia is streptococcus pneumonia, which can be prevented with the [pneumococcal vaccine](#). There are two [types of pneumococcal vaccines](#): [pneumococcal conjugate vaccine](#) (PCV) and pneumococcal polysaccharide vaccine (PPV). Both protect against invasive pneumococcal disease (such as meningitis and the blood infection referred to as septicemia), and the conjugate vaccine is proven to reduce the risk of pneumonia.

The [government funds](#) influenza (annually) and pneumococcal vaccines for people aged 65 and over.

Shingles is a reactivation of the chickenpox virus. It causes a high burden of disease in older people (who have had chickenpox before) and can lead to debilitating and chronic pain. The [shingles vaccine](#) is recommended for people aged 60 and over. The government funds it for people aged 70 to 79.

Australian travellers

Travel is a major vector for transmission of infections around the world, and travellers are at high risk of preventable infections. Most [epidemics of measles](#), for example, are imported through travel. People may be under-vaccinated for measles if they missed a dose in childhood.

Anyone travelling should discuss vaccines with their doctor. If unsure of measles vaccination status, vaccination is recommended. This will depend on where people are travelling, and may include vaccination for yellow fever, Japanese encephalitis, cholera, typhoid, hepatitis A or influenza.

Travellers who are visiting friends and relatives overseas often [fail to take precautions](#) such as vaccination and do not perceive themselves as being at risk. In fact, they are at higher risk of preventable infections because they may be staying in traditional communities rather than hotels, and can be exposed to risks such as contaminated water, food or mosquitoes.

Aboriginal Australians and Torres Strait Islanders

Indigenous Australians are at [increased risk of infections](#) and have access to funded vaccines against influenza (anyone over six months old) and pneumococcal disease (for infants, everyone over 50 years and those aged 15-49 with chronic diseases).

They are also advised to get hepatitis B vaccine if they haven't already received it. Unfortunately, overall [vaccine coverage for these groups](#) is low – between 13% and 50%, representing a real lost opportunity.

Migrants and refugees

Migrants and refugees are at risk of vaccine-preventable infections because they [may be under-vaccinated](#) and come from countries with a high incidence of infection. There is no systematic means for GPs to identify people at risk of under-vaccination, but the new [Australian Immunisation Register](#) will help if GPs can check the immunisation status of their patients.

The funding of catch-up vaccination has also been a major obstacle until now. In July 2017 the government announced [free catch-up vaccinations](#) for children aged 10-19 and for all newly arrived refugees. This covers any childhood [vaccine](#) on the [National Immunisation Schedule](#) that has been missed.

While this does not cover all under-vaccinated refugees, it is a welcome development. If you are not newly arrived but a migrant or refugee, check with your doctor about catch-up vaccination.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by University of New South Wales

Citation: Which vaccinations should I get as an adult? (2017, August 29) retrieved 17 April 2024 from <https://medicalxpress.com/news/2017-08-vaccinations-adult.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.