

Does alcohol affect the risk of developing an abdominal aortic aneurysm?

September 7 2017

In an analysis of published studies, lower levels of alcohol consumption were associated with a lower risk of abdominal aortic aneurysm until approximately 15 to 20 g/day, with an increasing risk thereafter. In the *British Journal of Surgery* analysis, the increase in risk beyond 2 units/day was stronger in men than in women.

An abdominal aortic [aneurysm](#) occurs when an area of the aorta—the main blood vessel that supplies blood to the abdomen, pelvis, and legs—becomes very large or balloons out. There are well established [risk factors](#) for the development of abdominal aortic aneurysm—such as increasing age, male sex, and smoking—but the role of alcohol remains uncertain.

The [analysis](#) combined results from 11 large prospective cohort studies, including more than 3500 cases identified from nearly 500,000 participants, followed-up for between 5 and 34 years.

"The association is fairly weak and the evidence is not strong for alcohol either decreasing or increasing the risk of abdominal aortic aneurysm by very much," said Dr. Darren Greenwood, senior author of the study. "If people want to stay healthy and active for longer, the biggest improvement they can make is to quit smoking."

More information: *British Journal of Surgery* (2017). [DOI: 10.1002/bjs.10784](https://doi.org/10.1002/bjs.10784)

Provided by Wiley

Citation: Does alcohol affect the risk of developing an abdominal aortic aneurysm? (2017, September 7) retrieved 6 May 2024 from <https://medicalxpress.com/news/2017-09-alcohol-affect-abdominal-aortic-aneurysm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.