

Chronic pain common in people living with HIV

September 14 2017

Because ongoing pain is a significant problem that affects 39 to 85 percent of people living with HIV, everyone with the infection should be assessed for chronic pain, recommend guidelines released by the HIV Medicine Association (HIVMA) of the Infectious Diseases Society of America (IDSA) and published in the journal *Clinical Infectious Diseases*. Those who screen positive should be offered a variety of options for managing pain, starting with non-drug treatment such as cognitive behavioral therapy, yoga and physical therapy, suggest the first comprehensive guidelines on HIV and chronic pain.

"Because HIV clinicians typically are not experts in [pain management](#), they should work closely with others, such as pain specialists, psychiatrists and physical therapists to help alleviate their patients' pain," said Douglas Bruce, MD, MA, MS, lead author of the guidelines, chief of medicine at Cornell Scott-Hill Health Center, and associate clinical professor of medicine at Yale University, New Haven, Conn. "These comprehensive guidelines provide the tools and resources HIV specialists need to treat these often-complex patients, many of whom struggle with depression, substance use disorders, and have other health conditions such as diabetes."

The guidelines recommend all people with HIV be screened for [chronic pain](#) using a few simple questions:

- How much bodily pain have you had during the week?
- Do you have bodily pain that has lasted more than three months?

Those that screen positive should undergo comprehensive evaluation, including a physical exam, psychosocial evaluation and diagnostic testing. Nearly half of chronic pain in people with HIV is neuropathic (nerve pain), likely due to inflammation or injury to the central or peripheral nervous system caused by the infection. Non-neuropathic pain typically is musculoskeletal, such as [low-back pain](#) and osteoarthritis in the joints.

"It has been long known that patients with HIV/AIDS are at high risk for pain, and for having their pain inadequately diagnosed and treated," said Peter Selwyn, MD, MPH, co-chair of the guidelines and professor and chair of the Department of Family and Social Medicine, and director of the Palliative Care Program for Montefiore Medical Center, Albert Einstein College of Medicine, Bronx, NY. "This is an aging population and the changing clinical manifestations of HIV, complexity of the disease and additional challenges related to substance abuse make treatment complicated. These guidelines help provide clarity in treating these patients."

HIV specialists should work with an interdisciplinary team to offer multi-modal treatment. The guidelines recommend offering alternative, non-pharmacological therapies first, including [cognitive behavioral therapy](#), yoga, physical and occupational therapy, hypnosis and acupuncture. If medication is needed, the guidelines recommend beginning with non-opioids, such as gabapentin (anti-seizure [medicine](#)) and capsaicin (topical pain reliever made from chili peppers), both of which help with nerve [pain](#).

"Opioids are never first-line," said Dr. Bruce. "The guidelines always recommend the most effective treatment with the lowest risk."

The online version of the guidelines includes an extensive list of resources for physicians to reference to help them treat the patients

comprehensively.

Provided by Infectious Diseases Society of America

Citation: Chronic pain common in people living with HIV (2017, September 14) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2017-09-chronic-pain-common-people-hiv.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.