


# Sleep deprivation is an effective antidepressant for nearly half of depressed patients

September 19 2017

---



Credit: CC0 Public Domain

Sleep deprivation—typically administered in controlled, inpatient settings—rapidly reduces symptoms of depression in roughly half of

depression patients, according the first meta-analysis on the subject in nearly 30 years, from researchers at the Perelman School of Medicine at the University of Pennsylvania. Partial sleep deprivation (sleep for three to four hours followed by forced wakefulness for 20-21 hours) was equally as effective as total sleep deprivation (being deprived of sleep for 36 hours), and medication did not appear to significantly influence these results. The results are published today in the *Journal of Clinical Psychiatry*.

Although total sleep [deprivation](#) or partial sleep deprivation can produce clinical improvement in depression symptoms within 24 hours, antidepressants are the most common treatment for depression. Such drugs typically take weeks or longer to experience results, yet 16.7 percent of 242 million U.S. adults filled one or more prescriptions for psychiatric drugs in 2013. The findings of this meta-analysis hope to provide relief for the estimated 16.1 million adults who experienced a [major depressive episode](#) in 2014.

Previous studies have shown rapid antidepressant effects from sleep deprivation for roughly 40-60 percent of individuals, yet this response rate has not been analyzed to obtain a more precise percentage since 1990 despite more than 75 studies since then on the subject.

"More than 30 years since the discovery of the antidepressant effects of sleep deprivation, we still do not have an effective grasp on precisely how effective the treatment is and how to achieve the best clinical results," said study senior author Philip Gehrman, PhD, an associate professor of Psychiatry and member of the Penn Sleep Center, who also treats patients at the Cpl. Michael J. Crescenz VA Medical Center. "Our analysis precisely reports how effective sleep deprivation is and in which populations it should be administered."

Reviewing more than 2,000 studies, the team pulled data from a final

group of 66 studies executed over a 36 year period to determine how response may be affected by the type and timing of sleep deprivation performed (total vs early or late partial sleep deprivation), the clinical sample (having depressive or manic episodes, or a combination of both), medication status, and age and gender of the sample. They also explored how response to sleep deprivation may differ across studies according to how "response" is defined in each study.

"These studies in our analysis show that sleep deprivation is effective for many populations," said lead author Elaine Boland, PhD, a clinical associate and research psychologist at the Cpl. Michael J. Crescenz VA Medical Center. "Regardless of how the response was quantified, how the sleep deprivation was delivered, or the type of depression the subject was experiencing, we found a nearly equivalent [response](#) rate."

The authors note that further research is needed to identify precisely how sleep deprivation causes rapid and significant reductions in [depression](#) severity. Also, future studies are needed to include a more comprehensive assessment of potential predictors of treatment outcome to identify those patients most likely to benefit from [sleep deprivation](#).

**More information:** Elaine M. Boland et al, Meta-Analysis of the Antidepressant Effects of Acute Sleep Deprivation, *The Journal of Clinical Psychiatry* (2017). [DOI: 10.4088/JCP.16r11332](https://doi.org/10.4088/JCP.16r11332)

Provided by Perelman School of Medicine at the University of Pennsylvania

Citation: Sleep deprivation is an effective anti-depressant for nearly half of depressed patients (2017, September 19) retrieved 20 April 2024 from <https://medicalxpress.com/news/2017-09-deprivation-effective-anti-depressant-depressed->

[patients.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.