


# Effect of osteoporotic fractures similar to diabetes burden

September 23 2017

---



(HealthDay)—Osteoporotic fractures and complications of diabetes

similarly impact quality of life (QoL), according to a study published online Sept. 12 in the *Journal of Evaluation in Clinical Practice*.

Zoltán Vokó, D.Sc., from Eötvös Loránd University in Budapest, Hungary, and colleagues analyzed data from 840 patients with osteoporosis and 943 patients with diabetes to estimate the effect of [osteoporotic fractures](#) and microvascular and macrovascular [complications](#) of diabetes on QoL using the EuroQol-5D (EQ-5D) questionnaire.

The researchers found that the effects of certain of osteoporotic fractures and diabetes complications were similar in size, as measured by the EQ-5D. Patients with hip fractures and compressions of the vertebrae experienced drops in QoL comparable to the most severe complications of diabetes, such as vision loss and amputations.

"Our comparative analysis, which showed that osteoporotic fractures reduce QoL as much as major complications of diabetes, highlights the need for comprehensive disease burden assessment, including losses in functionality and QoL, to support decision making," the authors write.

The study was funded by Novartis Hungary Ltd. and Sandoz Hungary.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Effect of osteoporotic fractures similar to diabetes burden (2017, September 23)  
retrieved 23 April 2024 from  
<https://medicalxpress.com/news/2017-09-effect-osteoporotic-fractures-similar-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.