


# Epileptic seizures show long-distance effects

September 19 2017, by Wiebe Van Der Veen

---



Measuring brain activity on micrometer and centimeter scales. Credit: University of Twente

The area in which an epileptic seizure starts in the brain, may be small but it reaches other parts of the brain at distances of over ten centimeters. That distant activity, in turn, influences the epileptic core, according to mathematicians and neurologists of the University of Twente and the university of Chicago.

The researchers describe these cross-connections in their paper in the *Proceedings of the National Academy of Sciences (PNAS)*. When a patient has a large, generalized epileptic seizure, often causing loss of consciousness, a large part of the [brain](#) is involved. But even in partial or

'focal' seizures, the activity isn't limited to the small area. In those cases, a patient can suddenly have difficulties speaking, show sudden movements or seem absent. The active area then is in the language centre or close to it, or in the motor cortex. As is shown now, even in these specific cases, a large part of the neocortex, the top layer of the brain, is involved. Advanced models of neuronal networks, together with [patient data](#), confirm this.

## Firing neurons

The [neurons](#) at the edge of the epileptic [core](#) fire in a violent way, as expected. This effect, however, could stay local, but in the sub millimeter range, the activity starts spreading. A wave front of very active neurons is developing. The mechanism of inhibition, by which the surrounding neurons should stop further spreading, clearly fails.

While the core of the seizure shows neurons firing at high frequencies, low frequency (6 Hertz) waves can be detected at larger distances in the neocortex. For the first time, mathematical models show that these two types of brain activity correlate. There are two areas that show interaction: the focal epileptic core causes activity at a distance in the neocortex, while the slow wave returning to the core, affects the firing in itself. What remains unclear is how the [seizure](#) stops, but the cross-scale interaction could well play a role in this.

## Patient data

In order to 'feed' their mathematical models with real life data, the researchers could use existing patient data gathered by the Columbia University Medical Center. There, a temporary implant was used for measuring local epileptic activity. This so-called 'Utah array', of 4 by 4 mm, consists of 96 microelectrodes. Using this, the activity of small

groups of neurons can be detected. In addition, activity on the centimeter scale was measured using an Electrocorticogram (ECoG). Other than an EEG, that measures through the skull, an ECoG is placed directly on the brain surface. These techniques are at the basis of epilepsy surgery, for localizing the area that is responsible for seizures and possibly remove it by surgery.

At the University of Twente, the research was done by the Nonlinear Analysis group of Prof Stephan van Gils and the Clinical Neurophysiology group of Prof Michel van Putten, who is a neurologist at the Medisch Spectrum Twente hospital in Enschede as well.

**More information:** Tahra L. Eissa et al. Cross-scale effects of neural interactions during human neocortical seizure activity, *Proceedings of the National Academy of Sciences* (2017). [DOI: 10.1073/pnas.1702490114](https://doi.org/10.1073/pnas.1702490114)

Provided by University of Twente

Citation: Epileptic seizures show long-distance effects (2017, September 19) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-09-epileptic-seizures-long-distance-effects.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--