

Researchers identify gene variants linked to a high-risk children's cancer

September 25 2017



Credit: CC0 Public Domain

Pediatric researchers investigating the childhood cancer neuroblastoma have identified common gene variants that raise the risk of an aggressive form of that disease. The discovery may assist doctors in better diagnosing subtypes of neuroblastoma.

Co-led by a genomics expert and a pediatric oncologist with expertise in <u>neuroblastoma</u>, a team from Children's Hospital of Philadelphia (CHOP)



reported that common variants in the *MMP20* gene are associated with deletions on chromosome 11q. "These inherited variants predispose to 11q deletions, which are a strong risk factor for an aggressive form of neuroblastoma," said co-study leader Hakon Hakonarson, MD, PhD, director of the Center for Applied Genomics at CHOP.

The study appeared online Sept. 18 in Nature Communications.

"We already knew that 11q deletions are biomarkers that predict poor outcomes in neuroblastoma," said co-study leader John M. Maris, MD, a <u>pediatric oncologist</u> at CHOP's Center for Childhood Cancer Research. "This new research helps us to more precisely predict how a neuroblastoma tumor will behave, so it improves our diagnostic capabilities."

A <u>cancer</u> of the peripheral nervous system that usually occurs as a solid <u>tumor</u> in a child's chest or abdomen, neuroblastoma is the most common cancer in infants. It accounts for a disproportionate share of cancer deaths in children.

The CHOP researchers previously identified six other genetic risk variants for neuroblastoma over the past decade, so the current investigation offers new insights into the genetic architecture of this complex form of cancer.

In the current study, the team performed a genome-wide association study in 113 tumors from neuroblastoma patients harboring 11q deletions, compared to 5,100 ancestry-matched controls. The research yielded common variants in the *MMP20* gene, located on chromosome 11q22.2. A replication study in 44 independent cases and 1,900 controls found similar results.

The authors added that further studies should investigate the mechanisms



by which MMP20 variants give rise to neuroblastoma tumors.

More information: Xiao Chang et al, Common variants in MMP20 at 11q22.2 predispose to 11q deletion and neuroblastoma risk, *Nature Communications* (2017). DOI: 10.1038/s41467-017-00408-8

Provided by Children's Hospital of Philadelphia

Citation: Researchers identify gene variants linked to a high-risk children's cancer (2017, September 25) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2017-09-gene-variants-linked-high-risk-children.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.