

Eating oily fish during pregnancy could prevent schizophrenia in the child, new study suggests

September 7 2017, by David Mazzocchi-Jones



Credit: CC0 Public Domain

Mice that are deprived of an essential fatty acid, called docosahexaenoic acid (DHA), during pregnancy, are more likely to produce pups that

display schizophrenia-like symptoms as adults, according to a [new study](#) from Japan.

DHA is an essential fatty acid – "essential" because our bodies can't produce it. It must be obtained from food. Oily fish, such as salmon and sardines, are good sources of dietary DHA. It is well understood that DHA plays a role in brain development. It is especially important during the last three months of pregnancy, and in the first two years of adolescence.

Studies have shown that babies fed on DHA-supplemented formula milk display higher visual acuity and [problem-solving at 10 to 12 months](#). In an animal study, rats deprived of DHA – resulting in a 50-80% reduction in DHA levels in the brain – were shown to have [impaired learning and memory](#). Conversely, dietary DHA supplementation has been shown to [improve learning and memory](#) in brain damaged lab rats.

In the Japanese study, conducted by researchers at the RIKEN Brain Science Institute in Tokyo, mice were fed on a [diet](#) free from DHA, prior to conception and up to the point the offspring had been weaned. The mouse pups were then given a standard diet, containing DHA, and tested at eight weeks, which roughly translates to human adolescence.

The cognitive function of the mice was assessed using standard mazes; they needed to find and remember the location of a food reward. And depression and motivation were assessed by monitoring the mice's general activity and how quickly they avoided open spaces in special mazes (mice prefer enclosed spaces).

Mice born from mothers fed on a diet that excluded DHA showed significantly lower performance on the range of tests, compared with mothers fed on a [standard diet](#). Consequently, these mice displayed schizophrenia-like symptoms including, impaired cognitive function, and

reduced motivation; characteristic of the early stage of disorder. This led the study's authors to suggest that getting enough DHA during pregnancy may protect against schizophrenia-like symptoms in the offspring.

The role of epigenetics

The mechanisms underlying how diet can play such an important role in brain function and health are poorly understood. The dogma of genetics being entirely based on what we inherit, rather than the environment we are exposed to, has been questioned by scientists in recent years. The advent in understanding of the concept of [epigenetics](#) has revolutionised the field of genetic science and provided a potential mechanism through which the environment exerts an influence on [genes](#).

Under epigenetic modification, certain [mechanisms](#) can change the way a gene functions, or is expressed – without changing DNA itself – resulting in vastly different outcomes. These mechanisms are in turn activated by environmental factors, including [diet](#).

In the Japanese study, the researchers investigated the levels of two genes ([Rxr and Ppar](#)), known to be associated with schizophrenia in humans. They found evidence that these genes had been modified by epigenetic factors, resulting in lower activity in the mice that displayed schizophrenia-like symptoms.

It's very difficult to draw a direct comparison between evidence gained from studies in [mice](#), to humans. However, the study identified similar low levels of the RxR and Ppar gene in hair follicle samples obtained from schizophrenic patients. This suggests that adequate levels of DHA in the maternal diet protects normal gene function, which in turn protects against expression of genes associated with schizophrenia.

Ultimately, given that [epigenetic modifications](#) to genes can also be

passed on to future offspring, this study provides further evidence for the critical role dietary levels of DHA play in brain function and health. Also, given that epigenetic modifications to genes can be passed on, adequate maternal nutrition is not just essential to their offspring, but also to future generations.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: Eating oily fish during pregnancy could prevent schizophrenia in the child, new study suggests (2017, September 7) retrieved 10 May 2024 from <https://medicalxpress.com/news/2017-09-oily-fish-pregnancy-schizophrenia-child.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
