

New study on the placebo effect and antidepressants in children and adolescents

September 15 2017

Although the clinical efficacy of antidepressants in children and adolescents is proven, it is frequently accompanied by side effects. In addition, the influence of the placebo effect on the efficacy of antidepressants is unclear. A meta-analysis of data from over 6,500 patients has now shown that, although antidepressants are more effective than placebos, the difference is minor and varies according to the type of mental disorder. The results were obtained by the University of Basel and Harvard Medical School and were published in the journal *JAMA Psychiatry*.

The most common mental [disorders](#) in children and adolescents include [anxiety disorders](#), depressive disorders, obsessive-compulsive disorder, and [posttraumatic stress disorder](#). Children and adolescents are often treated with psychotherapy together with newer [antidepressants](#) such as selective serotonin reuptake inhibitors (SSRIs) and [serotonin-norepinephrine reuptake inhibitors](#) (SNRIs).

Antidepressants work better than placebos but have more side effects

Psychologists from the University of Basel analyzed 36 drug trials in cooperation with colleagues from Harvard Medical School and the American National Institute of Mental Health. The trials included data from 6,778 children and adolescents up to the age of 18.

The results of the meta-analysis show that, although antidepressants work significantly better than placebos across the range of disorders, the difference is small and varies according to the type of [mental disorder](#). However, the results also showed that the [placebo effect](#) played a significant role in the efficacy of antidepressants. The study also found that patients treated with antidepressants complained of greater side effects than those who received a placebo. The side effects included everything from mild symptoms such as headaches to suicidal behavior.

Placebo effect stronger in cases of depression

According to the study, the effects of antidepressants and placebos vary according to the type of mental disorder: antidepressants have a greater specific effect in the case of anxiety disorders than depressive disorders. On the other hand, placebos have a stronger effect in depressed patients than in those with an anxiety disorder. Lead authors Dr. Cosima Locher and Helen Koechlin from the Division of Clinical Psychology & Psychotherapy at the University of Basel's Faculty of Psychology see potential here for new treatment concepts. These would make use of factors contributing to the placebo effect, applying them specifically to the treatment of depression.

Individual clarification needed

However, the meta-analysis also shows that antidepressants play an important role in the treatment of mental disorders in childhood and adolescence. "With this in mind, it's important to examine the relationship between clinical benefit and possible side effects in consultation with the doctor treating the patient," says Locher.

More information: Cosima Locher et al. Efficacy and Safety of Selective Serotonin Reuptake Inhibitors, Serotonin-Norepinephrine

Reuptake Inhibitors, and Placebo for Common Psychiatric Disorders Among Children and Adolescents, *JAMA Psychiatry* (2017). DOI: [10.1001/jamapsychiatry.2017.2432](https://doi.org/10.1001/jamapsychiatry.2017.2432)

Provided by University of Basel

Citation: New study on the placebo effect and antidepressants in children and adolescents (2017, September 15) retrieved 20 March 2024 from <https://medicalxpress.com/news/2017-09-placebo-effect-antidepressants-children-adolescents.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--