

Heat a trigger for seizures

September 7 2017, by Sanjay Mishra



Generalized 3 Hz spike and wave discharges in a child with childhood absence epilepsy. Credit: Wikipedia.

Fever is the most common trigger for seizures in children between 5 months and 6 years of age. But the underlying cause is not always clear.

Now in a study published in the journal *Epilepsy Research*, Jing-Qiong (Katty) Kang, M.D., Ph.D., and colleagues show in a genetically altered mouse model that elevated body temperature alone can increase vulnerability to [febrile seizures](#) even in the absence of infection or inflammation.

The scientists studied a [mouse model](#) of Dravet syndrome, a severe epilepsy disorder that has been linked to mutations in the GABRG2 gene. They previously reported an association between GABRG2 mutations and febrile seizures and related epilepsy spectrum.

Regardless of whether inflammation was present in the mice, a brief heating of the body increased brain excitation and caused a rapid rise in the body temperature and precipitated seizures in mice.

This finding suggests that genetic mutations that alter the brain's thermal control mechanism may trigger febrile seizures when body temperature is raised.

More information: Timothy A. Warner et al. Heat induced temperature dysregulation and seizures in Dravet Syndrome/GEFS+ Gabrg2+/Q390X mice, *Epilepsy Research* (2017). [DOI: 10.1016/j.epilepsyres.2017.04.020](#)

Provided by Vanderbilt University

Citation: Heat a trigger for seizures (2017, September 7) retrieved 11 May 2024 from <https://medicalxpress.com/news/2017-09-trigger-seizures.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.