

Weight loss after obesity surgery may reverse eye damage

September 13 2017



Credit: CC0 Public Domain

Early eye damage caused by obesity, high blood pressure and diabetes can potentially be reversed by obesity surgery, concludes a study presented at this year's Annual Meeting of the European Association for

the Study of Diabetes (EASD) in Lisbon, Portugal (11-15 September).

Changes to the vascular structure of the retina can reflect damage caused by [obesity](#), hypertension, diabetes, and a range of other chronic disease processes. Despite such diseases being commonplace in the population, the impact of weight loss resulting from [bariatric surgery](#) on the retinal microvasculature is not well known.

A research team led by Associate Professor Robyn Tapp of the University of Melbourne, Australia, and Leeds Beckett University in Leeds, UK, together with Dr Antti Viljanen (Turku University, Finland) and colleagues from Finland, Singapore, Hong Kong, and London studied the effects of weight loss on the retinal microvasculature of obese subjects following bariatric (obesity) surgery.

Eye changes were analysed from the SleevePET study conducted at Turku PET Centre, Finland. Twenty-two obese subjects scheduled for [obesity surgery](#) were recruited along with 15 lean, age-matched controls. Detailed eye examinations were performed at the start of the study and 6 months later to look for signs of obesity-related impairments to the vascular structure of the retina. These include narrowing of the arterioles which carry blood from the arteries to the capillary beds, and widening of the venules which return the blood from the capillaries to the larger veins.

The team found that in the 6 months following bariatric surgery, the obese subjects lost an average of 26kg while also showing improvements in the microvasculature of their retinas. Arteriolar narrowing and venular widening were both less pronounced, whereas no such changes took place in the control group.

The authors conclude that: "the findings suggest obesity-related microvascular changes are reversible after bariatric surgery-induced

weight loss. The capacity for the retinal microvasculature to improve following bariatric surgery suggests plasticity of the human microvasculature early in the disease course."

Provided by Diabetologia

Citation: Weight loss after obesity surgery may reverse eye damage (2017, September 13)
retrieved 4 May 2024 from
<https://medicalxpress.com/news/2017-09-weight-loss-obesity-surgery-reverse.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--