

Acetaminophen in pregnancy tied to ADHD risk in kids

October 30 2017, by Amy Norton, Healthday Reporter



(HealthDay)—Acetaminophen is considered the go-to pain medication

during pregnancy. But a new study adds to evidence linking the drug to an increased risk of behavioral issues in kids.

Researchers in Norway found that among nearly 113,000 children, those whose mothers used acetaminophen during pregnancy were slightly more likely to be diagnosed with attention deficit hyperactivity disorder (ADHD).

The link was, however, confined to longer-term use—particularly a month or longer.

When moms used acetaminophen for 29 days or more during pregnancy, their kids were twice as likely to be diagnosed with ADHD, versus women who did not use the [drug](#).

On the other hand, when expectant moms used the drug for a week or less, their kids showed a slightly decreased risk of ADHD.

Acetaminophen is best known by the brand name Tylenol, but it's an active ingredient in many pain relievers.

The new study, led by researcher Eivind Ystrom from the Norwegian Institute of Public Health in Oslo, is not the first to suggest a connection between prenatal acetaminophen and ADHD.

But experts said it's still hard to definitively pin the blame on the medication.

"That's the dilemma," said Christina Chambers, co-director of the Center for Better Beginnings at the University of California, San Diego.

Around half of pregnant women use acetaminophen at some point, so it's important to understand any risks, according to Chambers, who was not

involved in the study.

But with a study like this, she explained, it's difficult to know whether factors other than acetaminophen are to blame—including the underlying conditions the women had.

According to the study researchers, longer-term use was tied to ADHD whether women used it for pain, fevers or infections.

But if a woman was using the medications for weeks to treat a fever or infection, that indicates she was quite ill, Chambers pointed out.

And if she took the drug for chronic pain, Chambers said, that raises the question of what effects the pain condition could have on her pregnancy.

For now, Chambers stressed that pregnant women should not be scared off from using acetaminophen for a fever—since an untreated fever could carry risks.

"The last thing we'd want, heading into flu season, is for women not to use acetaminophen to get a fever down," she said.

"This study," Chambers added, "suggests that if there is a causal association between acetaminophen and ADHD, it's with more-chronic use."

Overall, more than 2,200 children in the study were diagnosed with ADHD—or about 2 percent of the whole group. The risk was just over twofold higher among kids whose mothers had used acetaminophen for 29 days or more during pregnancy.

Why would the medication affect ADHD risk? There are potential "biologically plausible" explanations, Chambers said.

The drug might, for instance, interfere with maternal hormones that are important for fetal brain development.

But even if long-term acetaminophen does influence ADHD development, Chambers said, this study suggests it's a "modest" effect.

"The [risk](#) to any one woman would be small," she said.

That said, Chambers pointed to a bigger-picture issue: Very few drugs have actually been studied in pregnant women, and fairly little is known about the safety of using any medication prenatally.

The study was published online Oct. 30 in the journal *Pediatrics*.

Dr. Mark Wolraich, a professor of pediatrics at the University of Oklahoma Health Sciences Center, wrote an editorial accompanying the study.

He agreed that the study points only to an association between acetaminophen and ADHD, which might reflect a "third factor" at play, such as the underlying condition that caused the women to take the drug.

Plus, Wolraich explained, many factors might affect the development of ADHD. The evidence is "much stronger" for a familial influence, since the disorder tends to run in families, he noted.

Still, Wolraich said, [pregnant women](#) may want to be "overly cautious" about using [acetaminophen](#) for any extended time. He suggested that [women](#) talk their doctor before using any medications.

More information: Christina Chambers, Ph.D., M.P.H., professor, department of pediatrics, and co-director, Center for Better Beginnings, University of California, San Diego; Mark Wolraich, M.D., professor,

pediatrics, Child Study Center, University of Oklahoma Health Sciences Center, Oklahoma City; Oct. 30, 2017, *Pediatrics*, online

The nonprofit Organization of Teratology Information Specialists has more on [medication use during pregnancy](#).

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Acetaminophen in pregnancy tied to ADHD risk in kids (2017, October 30) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2017-10-acetaminophen-pregnancy-tied-adhd-kids.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
