

Clinical evidence synopsis published for T2DM

October 18 2017



(HealthDay)—Adding a sulfonylurea or metformin to insulin is



associated with approximately a 1 percent reduction in hemoglobin A1c (HbA1c), and addition of a sulfonylurea (but not metformin) is associated with more hypoglycemic events, according a clinical evidence synopsis published online Oct. 17 in the *Journal of the American Medical Association*.

Rimke C. Vos, Ph.D., and Guy E.H.M. Rutten, M.D., Ph.D., from University Medical Center Utrecht in the Netherlands, summarized a research review of studies that examined the effect of the addition of oral hypoglycemic agents among patients with type 2 diabetes who do not achieve optimal glycemic control with insulin monotherapy.

The researchers found that adding a sulfonylurea to insulin was correlated with more hypoglycemic events than insulin alone; this was not seen for metformin. There was an approximate 1 percent decrease in HbA1c with the addition of a sulfonylurea or <u>metformin</u> to insulin.

"Future <u>randomized clinical trials</u> could address the effect of adding oral hypoglycemic agents to insulin therapy for the outcomes of diabetes-related morbidity and mortality, all-cause mortality, and quality of life," the authors write.

The authors disclosed ties to the pharmaceutical industry.

More information: <u>Abstract/Full Text (subscription or payment may</u> <u>be required)</u>

Copyright © 2017 HealthDay. All rights reserved.

Citation: Clinical evidence synopsis published for T2DM (2017, October 18) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2017-10-clinical-evidence-synopsis-published-t2dm.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.