

Critical care nurses should be prepared for open abdomen

October 11 2017



(HealthDay)—Critical care nurses need to be prepared to manage



patients with open abdomen, according to a report published in the October issue of *Critical Care Nurse*.

Eleanor R. Fitzpatrick, R.N., from the Thomas Jefferson University Hospital in Philadelphia, reviews case studies illustrating the implementation of the open abdomen technique and patient management strategies.

Fitzpatrick notes that the open abdomen technique and temporary abdominal closure after damage control surgery is rapidly becoming the standard of care for managing intra-abdominal bleeding and infectious or ischemic processes in <u>critically ill patients</u>. The most common indication for the open abdomen is damage control surgery for abdominal trauma, but this technique is being used in <u>patients</u> with abdominal compartment syndrome (ACS) due to acute pancreatitis and other disorders. Following the open abdomen <u>technique</u> and temporary abdominal closure, management priorities include hemodynamic stabilization, supporting respiratory function, preventing infection, managing pain, and monitoring for ACS.

"Determining the need for this potentially lifesaving intervention and managing the wound after the open <u>abdomen</u> has been created are all within the realm of critical care nurses," Fitzpatrick writes.

More information: Abstract

Full Text

Copyright © 2017 HealthDay. All rights reserved.

Citation: Critical care nurses should be prepared for open abdomen (2017, October 11) retrieved 6 May 2024 from https://medicalxpress.com/news/2017-10-critical-nurses-abdomen.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.