

## Type 1 diabetes and the microbiota—MAIT cells as biomarkers and new therapeutic targets

October 10 2017, by Véronique Etienne

Together with colleagues from AP-HP Necker–Enfants Malades Hospital in Paris, scientists from the Cochin Institute (CNRS / INSERM / Paris Descartes University) have discovered that the onset of type 1 diabetes is preceded by modification of MAIT lymphocytes. These cells—associated with mucosae and able to recognize elements of the microbiota—could therefore serve as new biomarkers for early detection and prevention of the illness. The researchers' findings are published in *Nature Immunology*.

Type 1 diabetes (T1D) is an autoimmune disease that develops when the immune system attacks <u>beta cells</u> in the pancreas. This destructive process is often attributed to autoreactive T lymphocytes. Yet the innate immune system, our body's first line of defense, also plays an important role in this disease. MAIT <u>cells</u> associated with the mucosae are an integral part of this front-line immunity. They are activated by bacteria—especially those of the intestinal flora, or <u>gut microbiota</u>. The research team wondered what role MAIT cells played in the T1D-related disturbance of this flora and of gut mucosal homeostasis.

Investigations, led by Agnès Lehuen relied on T1D animal models studied by her team at the Cohin Institute as well as blood samples from patients under the care of Dr. Jacques Beltrand from the Necker–Enfants Malades Hospital. Findings for both the samples and the animal models showed that MAIT cells are altered before T1D



develops. When T1D is diagnosed in children, the number of MAIT cells in their blood is lower than in nondiabetic children. Migration of MAIT cells to inflamed pancreatic tissue in T1D patients could explain the discrepancy. This is a very reasonable assumption since the pancreatic population of MAIT cells swells in nonobese diabetic (NOD) mice. (NOD mice develop a form of diabetes very similar to the human disease.) Furthermore, experiments performed by the researchers on human cells in vitro and on NOD mice suggest MAIT cells are directly linked to the destruction of pancreatic beta cells.

In addition to changes in their concentration and location within the body, it appears that a functional defect in MAIT cells is linked to the modifications of the gut mucosa seen in T1D patients. Indeed, a normal function of MAIT cells is to maintain the homeostasis of this mucosa. This capacity is compromised in T1D, making the gut mucosa permeable to bacteria, which, in turn, favors an autoimmune response. Ongoing research seeks to better understand the links between MAIT cells and the gut microbiota.

The team's discovery could lead to the development of new strategies for T1D treatment. Most importantly, we now know that MAIT cells are early biomarkers for this form of diabetes because they undergo changes before the disease develops. Their use as such should improve T1D prevention.

**More information:** Ophélie Rouxel et al. Cytotoxic and regulatory roles of mucosal-associated invariant T cells in type 1 diabetes, *Nature Immunology* (2017). DOI: 10.1038/ni.3854

Provided by CNRS



Citation: Type 1 diabetes and the microbiota—MAIT cells as biomarkers and new therapeutic targets (2017, October 10) retrieved 4 May 2024 from https://medicalxpress.com/news/2017-10-diabetes-microbiotamait-cells-biomarkerstherapeutic.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.