

# Study identifies risk and protective factors for depressive symptoms in African-American men

October 25 2017

---

African-American men report an average of eight depressive symptoms in a month, with family support, mastery, self-esteem, chronic stressors and discrimination among the factors that are significant to their psychological health, according to a new study led by researchers at Georgia State University.

Although African-Americans are less likely than whites to meet the criteria for major depressive disorder, they are at increased risk for depressive symptoms. Few studies have focused on identifying the risk and protective factors that contribute to depressive symptoms in African-American men, which this study addresses.

The researchers determined the stress process model, a framework for understanding health and health inequalities, was useful for identifying psychosocial risk and protective factors in African-American men, explaining about half (50 percent) of the depressive symptoms. The findings could be beneficial for directing health initiatives and policies aimed at improving the psychological health of this population.

They also found some of the risk and [protective factors](#) influence each other. For instance, self-esteem and mastery (how people perceive control over things that happen to them) play an important role in mitigating the negative psychological harm associated with lower-income neighborhoods. Family support also was a buffer for the harmful

[mental health](#) effects of [stress exposure](#). The increased depressive symptoms associated with higher levels of chronic stressors and daily discrimination are relatively lower among African-American men who report more family support.

The findings are published in a special issue on the Psycho-social Influences of African-Americans Men's Health in the *Journals of Gerontology: Psychological Sciences*.

"The factors that contribute to the mental health of African-American men are consistent with research on the factors that are important for the psychological well-being of the general population—coping resources, stress exposure and [economic conditions](#)," said Dr. Mathew Gayman, associate professor in the Department of Sociology at Georgia State. "However, African-American men report, on average, fewer coping resources, greater stress exposure and poorer economic conditions than the general population. It is the systematic disparities in these factors that contribute to race inequalities in psychological health. Ultimately, if we want to address the increased risk for mental health problems (and mental health generally) experienced by African-American men, we must address the social conditions and forces that shape race disparities in coping resources, stress exposure and economic conditions."

Using data from a community-based study of Miami-Dade County (Fla.) residents that was linked to neighborhood census data, the researchers surveyed nearly 2,000 people from different ethnic groups between 2000 and 2001. Analysis for this study was limited to only African-American men, a sample of 248 participants.

Depressive symptomatology was assessed using the Center for Epidemiologic Studies Depression scale. Participants were presented with statements such as "You felt depressed" and "You felt that you could not shake off the blues" in the past month and asked to give

responses ranging from 0 (not at all) to 3 (almost all the time). Higher scores represented more symptoms.

Various scales were also used to assess socioeconomic status (individual-level and neighborhood-level), social stressors, daily discrimination, perceived social support, mastery and self-esteem.

About 11 percent of the African-American men reported 16 or more depressive symptoms, a cutoff often used to estimate for clinical-level depression, although depressive symptoms in these men might be underreported because of gender differences in the expression of depression. Consistent with previous research, this study found individual socioeconomic status in African-American men was not associated with depressive symptoms, possibly because of the often-unrealized rewards associated with higher income and education among African-Americans.

However, the researchers determined African-American men living in lower socioeconomic neighborhoods experienced significantly more [depressive symptoms](#), highlighting the significance of neighborhood [socioeconomic status](#) in their [psychological health](#).

Because African-American men are more likely than white counterparts to live in lower-income neighborhoods, the researchers conclude that public health policies aimed at addressing poor mental health among African-Americans should account for neighborhood conditions. The findings also indicate that while self-reliance through mastery and self-esteem may be important for mitigating the psychological consequences associated with living in relatively poor neighborhoods, the ability to perceive support from one's family is important for minimizing the negative mental health consequences of stress exposure for African-American men.

Provided by Georgia State University

Citation: Study identifies risk and protective factors for depressive symptoms in African-American men (2017, October 25) retrieved 20 April 2024 from <https://medicalxpress.com/news/2017-10-factors-depressive-symptoms-african-american-men.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.