

Back on ice, but young hockey players' brains still recovering from concussion

October 25 2017

Hockey players in their early teens who have had a concussion may still have brain changes three months later, long after other symptoms have cleared and they are allowed to return to play, according to a study published in the October 25, 2017, online issue of *Neurology*, the medical journal of the American Academy of Neurology. The study looked at brain scans of boys who played in Bantam hockey leagues when body checking is first introduced.

"During the early teen years, the brain is still not fully developed and as it continues to grow and mature, it may be more vulnerable to brain injury," said study author Ravi S. Menon, PhD, of Western University in London, Ontario. "Our findings show the brains of young athletes may need more time to recover, which is important because without full recovery, players may be more susceptible to a second [concussion](#) that could cause life-threatening brain swelling and bleeding."

The study looked at male hockey players, ages 11 to 14 during a six-month hockey season. On average, they practiced twice a week and played at least one game a week. During the season, 17 players were diagnosed with a concussion, most often from falling and hitting the back of the head. Researchers then compared them to 26 healthy players, who had either never had a concussion or had not had one within at least six months.

At the beginning of the study, each player was given thinking, memory, and balance tests. Each player also had a [magnetic resonance imaging](#)

(MRI) scan of his brain. Those who were diagnosed with a concussion were given an immediate post-concussion assessment with MRI, and 14 players had a follow-up assessment with MRI three months later.

For all players with concussion, scores on the thinking and memory tests returned to normal before the three-month mark. On average, they required 24 days (ranging from 10 to 46 days) to recover and be cleared to return to play.

Compared to the [brain scans](#) of healthy players, the scans of the concussed players three months after the concussion showed signs of widespread damage to the white matter of the brain, as well as a 10 percent reduction in molecules associated with metabolism. The scans also showed other areas of the brain trying to create new connections in a possible attempt to recover function.

"More research with MRI is needed to further evaluate our findings because it is critical to understand how and when the adolescent human brain reacts and recovers from concussion," Menon said. "Current thinking, memory and balance testing may not be sensitive enough. These players were back on the ice when our study suggests their brains still needed time to heal."

One limitation of this study is that players were cleared to play at different times. Future MRI studies should follow [players](#) for a longer period of time to see how long it takes for the [brain](#) to heal.

Provided by American Academy of Neurology

Citation: Back on ice, but young hockey players' brains still recovering from concussion (2017, October 25) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-10-ice-young-hockey-players-brains.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.