

Ischemic stroke patients not receiving life-saving treatment, study finds

October 5 2017, by Jennifer Rainey Marquez



Credit: Georgia State University

Ischemic stroke patients who do not receive intravenous (IV) alteplase, a clot-dissolving medication, are significantly less likely to survive,

according to researchers at Georgia State University.

Ischemic [stroke](#) is the most common type of stroke. It occurs when a vessel that supplies oxygen-rich blood to the brain becomes blocked, often by a blood clot. IV alteplase was approved by the Food and Drug Administration as a [treatment](#) for acute [ischemic stroke](#) in 1996 and is known to reduce disability and improve functionality by restoring blood flow to the brain. Yet two decades later, less than 10 percent of patients receive the treatment.

The study, published in the *American Journal of Emergency Medicine*, analyzed 2008-13 data from the Georgia Coverdell Acute Stroke Registry, and linked it to 2008–13 hospital discharge and 2008–14 death data in Georgia. The investigators found that one year after discharge, [acute ischemic stroke](#) patients who did not receive IV alteplase treatment had a 49 percent higher likelihood of death.

"Clinicians may be hesitant to administer IV alteplase because of concerns about the drug's complications, which can include bleeding," said Dr. Moges Ido, the study's lead author and a part-time instructor at Georgia State's School of Public Health. "But this study indicates that unless major contraindications are present, patients should be offered this treatment as a life-saving measure."

The study authors examined data from 9,620 patients who were treated at 48 hospitals. They excluded patients who weren't eligible to receive the treatment because of contraindications, such as a recent history of brain surgery. Only a quarter of the eligible patients received IV alteplase.

Previous randomized studies have shown some long-term mortality reduction for patients treated with IV alteplase but the results were not statistically significant. This study demonstrates that IV alteplase is

associated with reduced risk of death, and that eligible [patients](#) should be identified and treated swiftly.

More information: Moges S. Ido et al. The impact of intravenous alteplase on long-term patient survival: The Georgia Coverdell acute stroke registry's experience, *The American Journal of Emergency Medicine* (2017). [DOI: 10.1016/j.ajem.2017.07.092](https://doi.org/10.1016/j.ajem.2017.07.092)

Provided by Georgia State University

Citation: Ischemic stroke patients not receiving life-saving treatment, study finds (2017, October 5) retrieved 27 April 2024 from <https://medicalxpress.com/news/2017-10-ischemic-patients-life-saving-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--